

FACTORS INFLUENCING THE INCIDENCE OF *EIMERIA LEUCKARTI* IN HORSES

Tauseef-ur-Rehman ^{1*}, Zohaib Saeed ², Muhammad Arfan Zaman ³,
Arbab Sikandar ³ and Hafiz Muhammad Ali ¹

¹Department of Parasitology, Faculty of Veterinary and Animal Sciences, The Islamia University of Bahawalpur, Bahawalpur -63100, Pakistan; ²Department of Parasitology, University of Agriculture, Faisalabad, Pakistan
³College of Veterinary and Animal Sciences, Jhang, Pakistan

*Corresponding author: drtauseef@iub.edu.pk

ABSTRACT

A cross-sectional study was planned to determine the prevalence rate of *Eimeria* (*E.*) *leuckarti* in horses and factors influencing its occurrence in district TT Singh, Punjab, Pakistan. From April 2009 to March 2010, 484 equine feces were collected and adopted a two-stage cluster method for random selection of horses. Samples were subjected to the quantitative floatation method. Found 50.41% of sampled horses were positive for *E. leuckarti*, with prevalence reaching its peak in August (OR=1.156; $\chi^2=20.055$). The lowest occurrence was recorded in April, May, and June, coinciding with the lowest humidity and precipitation of the year in Pakistan. It can be inferred from these observations that the wet season seems to be favorable for the propagation of this disease. Foals (124/197; 62.94%; OR=0.422; $\chi^2=20.825$) were found to be more susceptible ($P<0.05$) to *E. leuckarti* in comparison to adults (120/287; 41.81%) while mares with the prevalence of 56.48% (196/347; OR=0.512; $\chi^2=13.265$) were found to be more susceptible ($P<0.05$) as compared to and male horses (48/137; 35.04%). Horses kept on the non-cemented floor, mixed farming with other animals, given food on the ground, and watered in ponds/common places were found to have higher rates of prevalence of *E. leuckarti* ($P<0.05$) in comparison to those kept on cemented/partially cemented floor, separate farming, food in the trough and tap water, respectively. It is inferred from the findings of this survey that disease can be controlled by improving management systems.

Keywords: *Eimeria Leuckarti*, Epidemiology, Equine, Pakistan

Article History (2020-1149) || Received: 19 Nov 2020 || Revised: 29 Jan 2021 || Accepted: 03 Mar 2021 || Published Online: 05 Jun 2021

©2021 ABR - All Rights Reserved

1. INTRODUCTION

Equines are used for multipurpose in developing world. About two third of equine diseases are comprised of gastrointestinal diseases. Coccidiosis is an important disease that lowers the performance of horses. Species of *Eimeria* responsible for coccidiosis in horses include *E. leuckarti*, *E. solipedum* and *E. uniungulsti* (Ghahfarrokhi et al. 2014). *Eimeria leuckarti* was first named after the name of Rudolf Leuckart (Dubey and Bauer 2018). Dubey and Bauer (2018) stated that *E. leuckarti* is the only species responsible for coccidiosis in horses. *Eimeria* is an intracellular parasite of phylum Apicomplexa of Kingdom Protista which infects intestine. *Eimeria leuckarti* enter epithelial cells of intestine and then they penetrate to lamina propria (Hirayama et al. 2002). Signs and symptoms of coccidiosis in horses include acute and chronic diarrhea, acute abdominal pain, colic and unthrifty (Dubey and Bauer 2018). Infection is higher in foals as compared to adults. Foals can get oocysts of *E. leuckarti* as early as first day after birth (Dubey and Bauer 2018).

There are various reports of coccidiosis in domestic animals of Pakistan (Rehman et al. 2012; Khan et al. 2013) but there is not enough data on coccidiosis in horses of Pakistan. Moreover, there is scarce information on epidemiological factors affecting incidence of coccidiosis in horse. As the drug resistance has been increasingly reported, research interests of scientific community have been shifted to alternative ways that may comprehend use of plants, determinants influencing prevalence, use of probiotics (Abu-Akkada and Awad 2015) and search of novel vaccine targets (Khan et al. 2016). Deep understanding of epidemiology of coccidiosis in horses may assist in effective control and prevention of disease.

2. MATERIALS AND METHODS

Fecal samples of horses were collected from whole district of Toba Tek Singh. Sampling technique followed for random sampling was two stage cluster sampling with primary units being union councils. Sample size was determined through following formulae (Thrusfield 2008).

As there was no information available about prevalence of disease in study area, expected prevalence was considered 50%. Using formula given below, samples were collected from 45 union councils out of total 82 union councils. Map grid method was used for selection of union councils. Number of primary clusters i.e., union councils were determined using following equation.

$$g = \frac{1.96^2 T_s V_c}{d^2 T_s - 1.96^2 P_{exp} (1 - P_{exp})}$$

A pilot survey was done in three councils to calculate between cluster variance. Number of sampled horses was calculated by following formula.

$$n = \frac{1.96^2 P_{exp} (1 - P_{exp})}{d^2}$$

A questionnaire comprising of dichotomous and multiple-choice questions was pre-designed and further refined after pilot survey for collection of information regarding risk factors influencing the incidence of disease. Information regarding following variables were collected.

Age: Sampling animals were divided into two categories of foals and adults for collection of feces. Animals of less than one year age were categorized as foals and older than four years as adults.

Sex: Feces of both sexes were collected. About fifteen grams of feces from 3 different fecal balls were collected for preservation in potassium dichromate (2.5%). Fecal samples were then carried to laboratory. Fecal samples were examined through floatation technique using saturated solution of zinc sulphate following method of Zajac and Conboy (2006) with slight modification. Oocysts were then identified by morphological features as described by Soulsby (2006). To determine the effect of relative humidity, rainfall and temperature on occurrence of disease, meteorological data was provided by Meteorological cell, Department of Crop Physiology, University of Agriculture, Faisalabad.

Mantel–Haenszel (MH) chi-square and Odds Ratio (OR) were used to analyze the effect of factors with paired characteristics. SAS was used to carry out all statistical procedures (SAS 2003).

3. RESULTS

In this study 484 samples were collected and examined for presence of *E. leuckarti*. *Eimeria leuckarti* was only species which was identified in horses of T.T. Singh. It was found that 50.41 percent of horses were positive for *Eimeria leuckarti*. Statistical analyses (OR & Chi-square) showed that prevalence of disease is significantly influenced by sex and age of horse, floor type, feeding system and farming system. Highest prevalence (74.36%) was found in month of August (rainy season) (Fig 1). Fig. 1 shows a relation between rain fall, temperature, relative humidity, and occurrence of *Eimeria*. Prevalence of *Eimeria* seems to be high after highest amount of precipitation. Strong negative correlation was found between prevalence of disease and age of horse.




Fig. 1: Prevalence of *Eimeria* in horses of Toba Tek Singh district, Pakistan during changes in rain fall (mm), temperature and relative humidity in the period April 2009 to March 2010.

Higher prevalence was recorded in foals (62.94%) in comparison to adults (41.81%; OR=0.422; $\chi^2=20.8252$) (Table 1; Fig. 2). Oocyst discharge in feces of foals was found to be significantly higher as compared to those in adults. Significantly lower oocysts per gram (OPG) was recorded in feces of adults ($\chi^2=11.3257$; OR=2.012) in comparison to foals. Lower prevalence was recorded in male horses (35.04%; $\chi^2=13.2653$; OR=0.512) in comparison to females (56.48%). No association was found between prevalence of disease and body condition of horse (P=0.2434).

Table 1: Multivariate logistic regression analysis of associated factors with risk of *Eimeria* infection in horse

Term	Odd's ratio	CI	P Value
Month	1.156	1.088-1.228	0.000
Age	0.422	0.279-0.637	0.000
Sex	0.512	0.325-0.805	0.004
Floor	0.487	0.315-0.755	0.001
Farming type	2.671	1.759-4.057	0.000
Feeding system	1.741	1.123-2.700	0.013
Body condition	0.482	0.314-0.741	0.001

Higher Odds of finding oocysts was recorded in horses kept singly (58.06%; $\chi^2=11.8863$; OR=2.671) in comparison to horses raised in mixed species farming system (42.37%) (Table 1; Fig. 3). Horses fed in troughs (42.61%) were found to be less susceptible to disease in comparison to those given feed on ground (62.17%; $\chi^2=9.5973$; OR=1.741) (Table 1; Fig. 3). Floor type was found to be strongly associated with disease in horses. Higher prevalence was found in horses raised at non-cemented floor (58.04%) in comparison to partially cemented / cemented floor type (41.92%, $\chi^2=12.5122$, OR=0.487) (Table 1; Fig. 3).




Fig. 2: Prevalence of *Eimeria* in horses in relation to age and sex in the Toba Tek Singh district, Pakistan.




Fig. 3: Prevalence of *Eimeria* in horses for various feeding, floor and farming systems as well as body condition of the animal in the Toba Tek Singh district, Pakistan.

4. DISCUSSION

Results of present survey first time depicted association of various determinants with the incidence of coccidiosis in horses at large scale. There are various reports of *E. leuckarti* from different parts of world showing prevalence from 0.3 to 80% (Colombia 0.3%, Kashmir 0.34%, Nigeria 1.2%, Romania 1.90%, Greece 4%, Turkey 4.5% & 5.88%, Finland 1% & 5.8%, Poland 6.7%, Iran 7.68%, Nigeria 28.57%, Brazil 40.4%, USA 41.6%, Germany 80%) (Beelitz et al. 1994; Lyons and Tolliver 2004; Studzińska et al. 2008; Pandit et al. 2008; Papazahariadou et al. 2009; Umur and Açıci 2009; Ehizibolo et al. 2012; Ionitã et al. 2013; Aromaa et al. 2018; Ola-Fadunsin et al. 2019; Ramírez-Hernández et al. 2019; Hautala et al. 2019; Gomes et al. 2019; Khamesipour et al. 2021). Wide range of percentage may be attributed to different techniques of fecal examination and age of sampled animals (Studzińska et al. 2008). Coccidiosis in horses of Europe due to other coccidian have been reported by Gutiérrez-Expósito et al. (2017).

Coccidiosis in horses is believed to be mainly concern of young ones. Ionitã et al. (2013) compared prevalence of *E. leuckarti* in foal and adult Romanian horses and found 15% of foals positive in comparison to 1.9% overall prevalence. Ghahfarrokhi et al. (2014) recorded similar results. Ola-Fadunsin et al. (2019) observed an increasing pattern of incidence of disease with increasing age. Dubey and Bauer (2018) reviewed epidemiology of coccidiosis in horses and found that prevalence is higher in foals in comparison to adult horses in all studies. Complex hormonal profile and more contact with young ones may be reason of greater prevalence in females. Ola-Fadunsin et al. (2019) observed higher prevalence of coccidiosis in mares than males. Highest prevalence was observed during months of highest precipitation. Dolenc (1966) reported highest prevalence during months of October, March and April.

It is inferred from findings of this survey that disease can be controlled by improving management systems. Less prevalence in mix farming in comparison to single species farming may be due to fact that *Eimeria* species are host specific. So other species dilute oocyst concentration which resulted in lowered prevalence. Higher prevalence of *E. leuckarti* in animals fed on ground is in line with recommendations of Radostits et al. (2009) to avoid ground feeding as it enhances the chances of contamination of feed with oocysts. Papazahariadou et al. (2009) reported lower parasitic infections in stabled horses as compared to grazing horses. Reduced prevalence was reported by Mitchell et al. (2012) in cattle farms where watering utensil are emptied and cleaned frequently. Aromaa et al. (2018) reported lower prevalence of *Eimeria* in horse sheds that were cleaned daily. Easy cleaning of cemented floor may be reasoned for less prevalence in animals kept at cemented floor. Also cemented floor dries earlier and low moisture contents results in death of oocysts.

Conclusion: Coccidiosis in horses can be prevented by improving hygienic conditions. Coccidiosis in horses can be prevented by adopting cemented floor, cleaning of farms and hygienic feeding system. Foals and mares should be given special consideration especially during pregnancy.

Author's Contribution: TR conceived and designed experiments. TR performed the experiments. TR, MAZ and AS performed statistical analyses of experimental data. TR prepared the draft of the manuscript. ZS and HMA edited the manuscript. All authors critically revised the manuscript and approved the final version.

ORCID

Tauseef-ur-Rehman <https://orcid.org/0000-0003-3925-8028>
Zohaib Saeed <https://orcid.org/0000-0001-5706-7183>
Muhammad Arfan Zaman <https://orcid.org/0000-0001-7281-4781>
Arbab Sikandar <https://orcid.org/0000-0001-7966-7729>
Hafiz Muhammad Ali <https://orcid.org/0000-0002-9885-5933>

REFERENCES

- Abu-Akkada SS and Awad AM, 2015. Protective effects of probiotics and prebiotics on *Eimeria tenella*-infected broiler chickens. *Pakistan Veterinary Journal* 35: 446-450.
- Aromaa M, Hautala K, Oksanen A, Sukura A and Näreaho A, 2018. Parasite infections and their risk factors in foals and young horses in Finland. *Veterinary Parasitology: Regional Studies and Reports* 12: 35-38. <https://doi.org/10.1016/j.vprsr.2018.01.006>
- Beelitz P, Rieder N and Gothe R, 1994. *Eimeria leuckarti* infections in foals and their mothers in upper Bavaria. *Tierarztl Prax* 22: 177-181.
- De Souza PNB, Bomfim TCB, Huber F, Abboud LCS and Gomes RS, 2009. Natural infection by *Cryptosporidium* sp., *Giardia* sp. and *Eimeria leuckarti* in three groups of equines with different handlings in Rio de Janeiro, Brazil. *Veterinary Parasitology* 160: 327-333. <https://doi.org/10.1016/j.vetpar.2008.10.103>

- Dolenc M, 1966. Contribution to the knowledge on the biology, pathogenicity and distribution of *Eimeria (Globidium) leuckarti* Reichenow, 1950 (in Slovenian Language). Vet. Med. Thesis. Biotehnske Fakultete, University, Ljubljana, pp: 1-53.
- Dubey JP and Bauer C, 2018. A review of *Eimeria* infections in horses and other equids. *Veterinary Parasitology* 256: 58–70. <https://doi.org/10.1016/j.vetpar.2018.04.010>
- Ehizibolo DO, Kamani J, Ehizibolo PO, Egwu KO, Dogo GI and Salami-shinaba JO, 2012. Prevalence and Significance of Parasites of Horses in Some States of Northern Nigeria. *Journal of Equine Science* 23: 1-4. <https://doi.org/10.1294/jes.23.1>
- Ghahfarrokhi EK, Ahmadi A, Shahraki SG and Azizi HR, 2014. *Eimeria leuckarti* (Flesch, 1883; Reichenow, 1940) from worker horses and donkeys of Shahrekord, Iran. *International Journal of Advanced Biological and Biomedical Research* 2: 1980-1984.
- Gomes FA, Jansen AM, Machado RZ, Jesus Pena HF, Fumagalli MJ, Silva A, BF Alves, Roque ALR and Figueiredo LTM, 2019. Serological evidence of arboviruses and coccidian infecting horses in the Amazonian region of Brazil. *PLoS ONE* 14: e0225895. <https://doi.org/10.1371/journal.pone.0225895>
- Gutiérrez-Expósito D, García-Bocanegra I, Howe DK, Arenas-Montes A, Yeargan MR, Ness SL, Ortega-Mora LM and Álvarez-García G, 2017. A sero-survey of selected cystogenic coccidia in Spanish equids: first detection of anti-Besnoitia spp. specific antibodies in Europe. *BMC Veterinary Research* 13: 128. <https://doi.org/10.1186/s12917-017-1046-z>
- Hautala K, Näreaho A, Kauppinen O, Nielsen MK, Sukura A and Rajala-Schultz PJ, 2019. Risk factors for equine intestinal parasite infections and reduced efficacy of pyrantel embonate against *Parascaris* sp. *Veterinary Parasitology* 273: 52-59. <https://doi.org/10.1016/j.vetpar.2019.08.004>
- Hirayama K, Okamoto M, Sako T, Kihara K, Okai K, Taharaguchi S, Yoshino T and Taniyama H, 2002. *Eimeria* organisms develop in the epithelial cells of equine small intestine. *Veterinary Pathology* 39: 505-508. <https://doi.org/10.1354/vp.39-4-505>
- Ionitã M, Buzatu MC, Enachescu V and Mitrea IR, 2013. Coprological prevalence and intensity of gastrointestinal parasites in horses in some Romanian studs: preliminary data. *Agro-Life Scientific Journal* 2: 207-212.
- Khan JM, El-Ashram S, Liu HB, Khan SH, Ayan A, Liu XY, Wang H, Tang XM, Suo X and Hassan MF, 2016. Transgenic *Eimeria mitis* expressing chicken IL-4 mediated decrease in pathogenicity compared to wild type *Eimeria mitis* strains in broiler chickens. *Pakistan Veterinary Journal* 36: 415-420.
- Khan MN, Rehman T, Sajid MS, Abbas RZ, Zaman MA, Sikandar A and Riaz M, 2013. Determinants influencing prevalence of coccidiosis in buffaloes, Pakistan. *Pakistan Veterinary Journal* 33: 287-290.
- Khamesipour F, Taktaz-Hafshejani T, Tebit KE, Razavi SM and Hosseini SR, 2021. Prevalence of endo- and ecto-parasites of equines in Iran: A systematic review. *Veterinary Medicine and Science* 7: 25-34. <https://doi.org/10.1002/vms3.321>
- Lyons ET and Tolliver SC, 2004. Prevalence of parasite eggs (*Strongyloides westeri*, *Parascaris equorum*, and strongyles) and oocysts (*Eimeria leuckarti*) in the feces of Thoroughbred foals on 14 farms in central Kentucky in 2003. *Parasitology Research* 92: 400-404. <https://doi.org/10.1007/s00436-003-1068-2>
- Mitchell ESE, Smith RP and Ellis-Iversen J, 2012. Husbandry risk factors associated with subclinical coccidiosis in young cattle. *Veterinary Journal* 193: 119-123. <https://doi.org/10.1016/j.tvjl.2011.09.017>
- Ola-Fadunsin SD, Daodu n OB, Hussain K, Ganiyu IA, Rabi M, Sanda IM, Adah AS, Adah AD and Aiyedun JO, 2019. Gastrointestinal parasites of horses (*Equus caballus* Linnaeus, 1758) and risk factors associated with equine coccidiosis in Kwara and Niger States, Nigeria. *Sokoto Journal of Veterinary Sciences* 17: 35-43. <https://doi.org/10.4314/sokjvs.v17i3.6>
- Pandit BA, Shahardar RA and Jeyabal L, 2008. Prevalence of gastrointestinal parasitic infestation in equines of Kashmir Valley. *Veterinary Scan* 3: 8-10.
- Papazahariadou M, Papadopoulos E, Diakou A and Ptochos S, 2009. Gastrointestinal Parasites of Stabled and Grazing Horses in Central and Northern Greece. *Journal of Equine Veterinary Science* 29: 233-236. <https://doi.org/10.1016/j.jevs.2009.03.005>
- Radostits OM, Blood DC and Gay CC, 2009. *Veterinary medicine. A textbook of the diseases of cattle, sheep, pigs, goats, and horses*. Bailliere Tindall, London, pp: 1181.
- Ramírez-Hernández A, Polo G, Robayo-Sánchez LN, Cruz-Maldonado OA, Imbacuán-Pantoja WO and Cortés-Vecino JA, 2019. Gastrointestinal and pulmonary parasites of working horses from Colombia. *Veterinary Parasitology: Regional Studies and Reports* 17: 1-6. <https://doi.org/10.1016/j.vprsr.2019.100296>
- Rehman T, Khan MN, Sajid MS, Iqbal Z, Javid MT, Riaz M and Ahmad M, 2012. Epidemiology of *Eimeria* and associated risk factors in goats of district Toba Tek Singh, Pakistan. *Indian Journal of Animal Science* 82: 1-4.
- SAS, 2003. SAS Statistical Software. Version 9.1. SAS Institute Inc., Cary, NC, USA.
- Soulsby EJJ, 2006. *Helminths, arthropods and protozoa of domesticated animals*. Baillier Tindall, London, pp: 289-326.
- Studzńska MB, Tomczuk K, Sadzikowski AB, 2008. Prevalence of *Eimeria leuckarti* in young horses and usefulness of some coproscopical methods for its detection. *Bulletin of the Veterinary Institute in Pulawy* 52: 541-544.
- Thrusfield M, 2008. *Veterinary epidemiology*. Blackwell, London, UK, pp: 39-64.
- Umur S and Açıci M, 2009. A survey on helminth infections of equines in the central black sea region, Turkey. *Turkish Journal of Veterinary and Animal Science* 33: 373-378. <https://doi.org/10.3906/vet-0712-6>
- Zajac AM and Conboy G., 2006. *Veterinary Clinical Parasitology*. Wiley-Blackwell, USA, pp: 114