

## UTILIZING CROP GENETIC RESOURCES IN MODERN BREEDING TO ACHIEVE SUSTAINABLE GOALS IN A CHANGING CLIMATE

Zahid Manzoor <sup>1,\*</sup>, Umbreen Shahzad<sup>2</sup>, Muhammad Sheeraz Qadir<sup>3</sup>, Muhammad Shah Jahan<sup>2</sup>, Zeshan Hassan<sup>2</sup>, Iqra Parveen<sup>2</sup>, Muhammad Shakeel Nawaz<sup>3</sup>, Ahsan Raza<sup>3</sup>, Zeba Shahnaz<sup>4</sup>, Memoona Bibi<sup>4</sup>, Uzma Aslam<sup>4</sup>, Iqra Khalid<sup>4</sup>, Shahid Nadeem<sup>4</sup>, Muhammad Yasir Khan<sup>4</sup> and Misbah Ghulam Rasool<sup>5</sup>

<sup>1</sup>National Key Laboratory of Crop Genetic Improvement, College of Plant Science and Technology, Huazhong Agricultural University, Wuhan, China

<sup>2</sup>College of Agriculture, University of Layyah, Pakistan

<sup>3</sup>Ayub Agriculture Research Institute, Vegetable Research Station Karor, Layyah, Pakistan

<sup>4</sup>Department of Plant Breeding and Genetics, Faculty of Agriculture, University of Agriculture, Faisalabad, Pakistan

<sup>5</sup>Department of Botany, University of Agriculture Faisalabad, Pakistan

\*Corresponding author: [zahidmanzoor041@gmail.com](mailto:zahidmanzoor041@gmail.com)

### ABSTRACT

Crop improvement has been both a vital goal and a challenge for humans since the advent of modern cultivation. Farmers and breeders have long sought to produce higher yields of food; therefore, they initially focused on selecting the most productive plants. They always adopted high-yielding crops and discarded the other types, resulting in a loss of genetic diversity. Developing breeding programs and utilizing crop genetic resources enabled the achievement of sustainable goals in food production and the maintenance of increased genetic diversity in various gene banks. Modern plant breeding has introduced methods to generate diversity in crop germplasm, and these newly formed types can also be conserved alongside parent lines, landraces, or wild types. In this review, we focus on discussing the utilization and conservation of crop genetic resources through the application of modern breeding tools and conservation strategies.

**Keywords:** Genetic resources, Modern breeding, Genetic diversity, Climate change, Sustainable production

---

Article History (ABR-25-027) || Received: 25 Mar 2025 || Revised: 29 Apr 2025 || Accepted: 15 May 2025 || Published Online: 16 Jun 2025

This is an open-access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

---

### 1. INTRODUCTION

Agriculture is an occupation in which more than 50% of the world's population is directly or indirectly engaged, especially in developing countries, where the sole income of many poor communities depends on agriculture. The hot topic nowadays is unstable environmental conditions, erratic rainfalls, global warming, and the most recent inclusion is that global cooling is a part of our lives (Hakeem, 2015). There is an increase of 1.0, 1.6, 0.9 and 1.3% in rice, maize, wheat, and soybean production per annum, respectively, which is less than 2.4%, to meet acute hunger and food demand until 2050 (Ray et al., 2013). Considering the current scenario, agricultural production cannot feed the world population, which is increasing so rapidly that it is expected to reach 10 billion in the coming years (Razzaq et al., 2021). This puts additional pressure on sustainable food production.

A growing threat crop faced during the past few decades is dynamic climatic conditions. This is one of the most significant uncontrollable factors affecting agricultural growth and production worldwide (Razzaq et al., 2021; Zafar et al., 2025). Local temperature and precipitation are vital to productivity and crop variety. Due to increased temperature, erratic rainfall, and fluctuations in precipitation, agricultural inputs and outputs vary severely across the globe (Kim, 2012). The farmer must adopt climate-smart agricultural practices in response to changing local environmental conditions, which require additional effort and resources.

Furthermore, the effect of these changes cannot be measured. The daunting fluctuations in climate directly and indirectly impact crop yield, irrigation/canal water availability. According to the report issued by the International Food Policy Research Institute (IFPRI, 2009), the report presented a detailed crop growth modeling study using two climate scenarios to simulate future climate conditions. The outcome suggested the negative impact of a changing climate on human well-being and crops (Nelson et al., 2009). Competing with these climatic factors is necessary to maximize agricultural outcomes. The only way is to create novel variability and a source population within the crop species that can withstand biotic and abiotic climatic effects.

Genetic diversity is a prerequisite to initiating any breeding program (Reddy et al., 2018). The presence of enough

---

**Citation:** Manzoor Z, Shahzad U, Qadir MS, Jahan MS, Hassan Z, Parveen I, Nawaz MS, Raza A, Shahnaz Z, Bibi M, Aslam U, Khalid I, Nadeem S, Khan MY and Rasool MG, 2025. Utilizing crop genetic resources in modern breeding to achieve sustainable goals in a changing climate. *Agrobiological Records* 20: 74-83. <https://doi.org/10.47278/journal.abr/2025.021>

genetic variability is the basis for present and future food security. It is essential to break yield barriers and introduce new cultivars to meet the ever-increasing demand for food among the growing population (Mashilo et al., 2017). The selection of desired traits requires a wider genetic base to induce a broad sense of heritability in germplasm in changing environmental conditions. DNA-based molecular markers play a crucial role in the early identification of genes that need to be incorporated to mitigate climatic stress (Awan et al., 2021; Razzaq et al., 2021). These variations can be achieved by collecting varieties, landraces and obsolete cultivars through the acclimatization of imported germplasm. If there is not enough variation present, manual efforts like induced mutation, polyploidy, genetic engineering and other modern techniques can serve as a helping hand to create novel variations (Awan et al., 2021).

The Latest research proposes that there is a 21% reduction in total factor productivity for worldwide agriculture since 1961 due to climatic trends (Ortiz-Bobea et al., 2021). Apart from conventional breeding, which needs more duration and further causes genetic erosion and loss of variability because of continuous selection (Abberton et al., 2016) modern breeding approaches like molecular breeding using DNA markers and genome editing tools (CRISPR Cas 9) (Razzaq et al., 2021). Along with innovative agricultural practices and improvement in plant microbiome conditions, it is an excellent way forward to attain desired food production goals. Next-generation breeding tools are a few methods to develop varieties resilient to erratic climatic conditions by utilizing genetic resources. Recent advances in genomic-assisted breeding (GAB) give a full landscape of genetic variability within a crop species to revive the lost gene. A well-known gene editing technique, and still evolving, known as next-generation CRISPR Cas-9 (clustered regularly interspaced short palindromic repeat), has revolutionized the idea that genome editing is restored for crop improvement (Jinek et al., 2012).

Furthermore, the discoveries of Cas orthologs, Cas9, Cas12, Cas13 and Cas14, has strengthened the editing efficacy (Koonin et al., 2017). We should move agriculture toward modernization and automation by adopting climate-smart agronomic practices and speed breeding involving artificial intelligence and machine learning. The amalgamation of genomic and phenomic tools with speed breeding allows us to quickly identify the desired gene. High-throughput phenotyping is a multidisciplinary approach to accelerate next-generation speed breeding and ultimately accelerate crop modification and improvement programs to develop climate-ready crops toward global food security (Fasoula et al., 2020).

This review highlights the use of genetic variability in conserving species in response to changing climatic trends. The conventional breeding approach undoubtedly introduces novel variability by crossing diverse populations, but it also causes genetic drag, genetic extinction, hybridization bottlenecks, and a laborious selection process. To meet the food demand of this ever-increasing population and achieve the zero hunger goals, we need to adopt modern breeding methods that utilize crop genetic diversity.

## 2. GENETIC RESOURCES TO CONSERVE BIODIVERSITY

Genetic Resources are essential in serving the provisioning ecosystem based on genetic material relevance to humans. They are also involved in improving and domesticating species that can be utilized in food production, drugs, fiber production, and many other food manufacturing for mankind (Nyadanu et al., 2017). Genetic resources also have some gradual and sudden natural and human-induced factors, for example, climate change and fire that can lead to a big loss. So the proper tools can be used to evaluate genetic resources such as genetic markers, golden trials, seed banks (Dreisigacker et al., 2005), pollen banks, and cryopreservation, and field gene banks (Pastorino & Marchelli, 2022). Advancements in conserving genetic resources for agricultural and food production are increasing daily and have expanded the scope of genetic resource conservation. For example, using economic tools (Wellband et al., 2021) such as gene banks, may be highly beneficial for managing genetic resources (Gollin, 2020). The functional food supply is based on plant biodiversity and benefits the resilience of climate change and ecosystem functions. To enhance the value of crop conservation, gene bank management and the integration of genomics and phenomics were employed, resulting in more sustainable and efficient resource conservation. For example, Genebanks and the integration of management tools may be involved in the management of genetic conservation data which is a major challenge in plant biodiversity that would be beneficial for breeders and other users (Ebert & Engels, 2020).

A decrease in biodiversity inhibits the capability of ecosystem functions to provide the sustainability of genetic resources and other services like hygienic food and water, and also decreases the genetic variability that may potentially reduce the resources required for future use. To overcome this reduction, some measures were used to protect biodiversity; for example, modified habitats, natural habitats, legally protected areas, and invasive alien species (de Boef et al., 2013).

### 2.1. Types of Conserving Biodiversity

The sustainability index of the environment is directly proportional to the conservation of biodiversity, and in this regard, the preservation of biodiversity should be the priority of all nations (Torchio et al., 2020). It is mandatory to have the centers of diversity information for fragile habitats. Furthermore, the conservation of biodiversity is the

obligation and responsibility of every nation for environmental sustainability (Arora, 2018).

The types of biodiversity conservation involve ex-situ and in-situ conservation (Mondal & Palit, 2022). The ex-situ strategy conserves all living organisms in an artificial habitat that considers natural living habitats, such as cryopreservation, Botanical gardens, aquariums, zoos, DNA banks, etc. (Kasso & Balakrishnan, 2013). In-situ conservation is the strategy to conserve endangered and wild species and refers to the restoration and protection of the population of species and their habitats (Wang & Li, 2021). The examples of in-situ conservation of biodiversity are elaborated in Table 1.

**Table 1: In-Situ Conservation of Plant Biodiversity**

In-situ Conservation	
Types	Examples
<b>Biosphere Reserves</b>	
These are the sites nominated by national governments that have multi-protected areas of an ecosystem open to researchers and the traditional lifestyles of inhabitants (Mengist et al., 2022).	Nanda Devi Sundarban Nokrek Manas
<b>National Parks</b>	
These are reserves for the conservation of environmental sustainability and wildlife, maintained by the government. Its limitations are highly demarcated, and humans, including habitat, Bandipur National Park cultivation, forestry, and grazing, are prohibited (Beissinger, 2022).	Kanha National Park
<b>Wildlife Sanctuaries</b>	
The type of conservation biodiversity that is involved in the regions where only wild animals and human activities such as cultivation, harvesting, forest products, and collection of wood are allowed (Rahman et al, 2017).	Abohar Wildlife Sanctuary Ghana Bird Sanctuary Mudumalai Wildlife Sanctuary

## 2.2. Modern Tools in Plant Breeding to Boost Biodiversity

Before the laws of genetics were known, the art of breeding had already been developed. By using the application of the principle of genetics to crops, a phenomenal increase in crop production occurs, particularly in cereal grains (Jeeterwal & Nehra, 2018). But conventional breeding methods may take 10 or more years to transfer a trait from one parent to another cultivar and it also has various limitations in boosting crop biodiversity.

To meet the future demand and boost crop biodiversity, one way of increasing agricultural production might be to breed with or without GMOs, as it requires less water, has better nutrient usage, and is adapted to climate change (Głowacka et al., 2018). But it is the only way to boost biodiversity and in the current situation, the yield enhancement in major crop species (varying from 0.8–1.2%) must be doubled (Li et al., 2018). So, to conserve crop biodiversity, scientists have adapted new techniques and are moving towards modern genetics and breeding of crop plants. Some biodiversity conservation applications have been mentioned that require whole-genome sequences. So, mostly used approaches is reduced representation sequencing (RRS) that targets a relatively large, about 1% unlinked, representative subsequence of the genome, reducing costs per sample and letting more depth of coverage per locus or larger number of individuals (Alex Buerkle & Gompert, 2013). Now in current studies, restriction site-associated DNA sequencing (RADseq) uses sequence fragments adjacent to sites cut by restriction enzymes (Andrews et al., 2016). RADseq use is not restricted to model organisms because it does not require prior genomic information, as microsatellites are obtained from genomic data. Thousands of genome-wide SNPs can be identified for the cost of developing some microsatellites so they are markers of choice (Kjeldsen et al., 2016). SNPs' main advantage is that they are directly comparable between labs for collaborative studies (De Groot et al., 2016). (Kim et al., 2016) reported the whole-genome sequencing of 137 rice mini core collections, which presented approximately 25,604 rice germplasms in the Korean genebank of the Rural Development Administration (RDA) based on the Nipponbare reference genome, and resequencing data yielded more than 15 million (M) SNPs and 1.3 M INDELs. Approximately 2,046,529 high-quality SNPs were assigned to the accessions of relevant subgroups for the study of rice mini core, utilizing phylogenetic and population analysis. These SNPs capture the evolutionary signatures present in rice subpopulations. Similarly, by using 201,817 SNPs, a population structure analysis of 300 rapeseed Chinese germplasm with 22 other accessions of different origins was carried out to obtain sequencing (Zhou et al., 2017).

For sequence messenger RNA (RNAseq), the next generation sequencing (NGS) can be used as it is the transcriptome (Todd et al., 2016). It requires a focus on only the functional parts of the genome, and RNA-seq requires high-quality tissue samples, but RNA is rapidly degraded. That is why its use in conservation studies is restricted. The NGS platforms can be used to assess DNA methylation, a major mechanism of epigenetic modification (Verhoeven et al., 2016). The difference can be identified in methylated and unmethylated cytosine bases by sodium bisulfite during sequencing (bsRADseq). Other epigenetic modifications, such as chromatin and histone, have a lack of interest in conservation as these are not heritable (Verhoeven et al., 2016).

**Citation:** Manzoor Z, Shahzad U, Qadir MS, Jahan MS, Hassan Z, Parveen I, Nawaz MS, Raza A, Shahnaz Z, Bibi M, Aslam U, Khalid I, Nadeem S, Khan MY and Rasool MG, 2025. Utilizing crop genetic resources in modern breeding to achieve sustainable goals in a changing climate. *Agrobiological Records* 20: 74-83. <https://doi.org/10.47278/journal.abr/2025.021>

Over the past two years, CRISPR/Cas9 has displaced other alternative tools such as zinc-finger nucleases and transcription-activator-like effector nucleases (TALENs). It is less cost-effective and simpler to use (Du et al., 2016). Performing in the molecular biology lab, genome editing with CRISPR/Cas9 is very cheap, easy and versatile, which is the main factor of biodiversity conservation. Some applications are currently being explored in crops (Weeks et al., 2016) and livestock (Reardon, 2016). That has relevance in conservation. These crops such as maize (Svitashev et al., 2015), wheat (Sánchez-León et al., 2018), rice (Zhou et al., 2015), tomato (Brooks et al., 2014), soybean (Han et al., 2019), cotton (Zhao et al., 2017) have improved by using different genetic resources. Using these techniques also contributes to the conservation of biodiversity.

In the future, it should become possible to obtain a complete understanding of the link between genotype, phenotype, and environment by omic technologies (Yugi et al., 2016). Although the applications of integrated omics have not yet been developed for conservation (Table 2).

**Table 2:** Recent studies on utilizing crop genetic resources in breeding and pre-breeding programs

Targeted Crop	Genetic Resource	Technology	Outcome	References
Wheat	32 genotypes	RFLP Markers	Assessment of genetic variability and pathogenicity through the identification of 30 unique and 36 shared alleles	Verma & Singh (2020)
Rice	Two rice cultivars (Yukinko-mai and Kaijin) from the Niigata Agricultural Research Institute's Crop Research Center and Iwate Biotechnology Research Center, Japan	SNP marker-assisted selection + speed breeding	Salt tolerance improvement	Rana et al. (2020)
Rice	SWARNA SubI, CR Dhan 800	Marker-assisted backcrossing	Submergence tolerance, bacterial blight resistance and high-yielding near-isogenic lines under changing climate	Mohapatra et al. (2021)
Rice	60 natural accessions	QTL-based- Marker-assisted selection	Identification of genotypes less responsive to Cd toxicity and boosting genotypic effects	Sun et al. (2022)
Tomato	63 accessions of 14 Spanish Tomato landraces	The specific combination of SNP markers	A high degree of variation in the population, heterozygosity detected spontaneous crossing, genetic fingerprinting of landraces and In-situ conservation of landraces was suggested	Cortes-Olmos et al. (2015)
Tomato	426 tomato accessions	High-density SNP genotyping	Identification of 7 sub-populations by Principal Component Analysis, linkage disequilibrium in each chromosome of these populations, different patterns of genetic variations,	Sim et al. (2012)
Tomato	142 FI cultivars of 4 different classes	SNPs were generated from genotyping by sequencing	Genome-wide SNP discovery, DNA barcoding, variety identification	Kim et al. (2021)
Cotton	50 representative Pakistani cotton cultivars	SSR primer pairs-based selection	Genetic diversity assessment of cotton genotypes by using SSR markers	Dahab et al. (2013)
Cotton	25 cotton genotypes	Selection based on SSR and ISSR markers	Genetic diversity determined by SSR and ISSR markers	Bardak & Bölek (2012)
Cotton	12 cotton varieties	RAPD primer series used for selection	Assessment of variability by using RAPD markers	Bukhari et al. (2021)
Millet	40 accessions of Italian millet	SSR markers developed through SSR- an enriched library from genomic DNA	Development and use of novel SSR markers for molecular genetic diversity	Zhao et al. (2012)
Chilli	20 local chili accessions of Bangladesh	Selection based on SSR markers	To assess the molecular genetic diversity by SSR markers	Sharmin et al. (2018)
Sugarcane	PSJT-941, PS-862, and BL varieties	In vitro selection using PEG-selecting agents	In-vitro selection of sugarcane putative mutant for drought stress	Hartati et al. (2021)
Common bean	IAC Formoso x BRS Perola	Marker-assisted backcrossing	Combining disease resistance and postharvest quality traits by early marker-assisted backcrossing in Carioca beans	Paulino et al. (2021)
Chickpea	ICCV 92944, ICCV 00108, ICC 4958 (Donor parent)	Marker-assisted backcrossing	Drought tolerance root traits into commercial chickpea varieties	Kosgei et al. (2022)
Wheat	4 cultivars	Backcross selection and marker-assisted selection	To improve stripe rust resistance by backcrossing and marker-assisted selection	Zhang et al. (2022)
Barley	Garnal x Baisheshak, Natali x Auksiniai	Zinc- Finger Transcription Factors and SNP markers	Use of ZFT Factors and SNP markers for drought tolerance	Baidyussen et al. (2021)

### 3. MARKER-ASSISTED BREEDING AND GENOME SEQUENCING

Advancements in genomic research are used to identify the genome sequence and pinpoint several genes precisely via linkage to DNA markers. For example, several genes are tightly linked to genetic markers that confer resistance to virus diseases, drought, salinity, bacterial blight, and submergence, as well as improve the quality of

**Citation:** Manzoor Z, Shahzad U, Qadir MS, Jahan MS, Hassan Z, Parveen I, Nawaz MS, Raza A, Shahnaz Z, Bibi M, Aslam U, Khalid I, Nadeem S, Khan MY and Rasool MG, 2025. Utilizing crop genetic resources in modern breeding to achieve sustainable goals in a changing climate. *Agrobiological Records* 20: 74-83. <https://doi.org/10.47278/journal.abr/2025.021>



traits (Duitama et al., 2015). Marker-assisted selection can monitor the presence or absence of these genes in breeding populations. This strategy may provide opportunities for high-yield, better-quality crop cultivars and stress-resistant cultivars for breeders. Studies have been conducted that utilize the positive aspects of this breeding device, which apply to new crop varieties and provide conservation genetic diversity (Lei et al., 2021; Munda et al., 2022). Marker-assisted selection also provides opportunities to develop high resistance to insects and diseases, which would benefit breeders (Haque et al., 2021; Jena & Mackill, 2008). Marker-assisted backcrossing effectively integrates major genes or traits with high effectiveness into different cultivated varieties (Jena & Mackill, 2008). A study highlights the use of marker-assisted backcrossing (MABC) to conserve the critical germplasm or genetic resources of rice crops under various types of flooding stress (Panda et al., 2021). Plant breeding is based on selecting phenotypes that require field observations, breeding cycles, and enormous resources (Michel et al., 2017). Recurrent selection is a breeding strategy based on population improvement that involves selecting phenotypes to enhance the frequency of favorable alleles between individuals through repeated intercrossing (Singh et al., 2021). A study conducted in China demonstrated this strategy for developing new premium varieties of the crop, which are characterized by high quality and disease resistance, and can be cultivated in monsoon climate conditions (Wang & Li, 2021).

The advancement of revolutions in genomics is caused by third-generation or long-read sequencing technologies. The main aim of this advancement is to study transcriptomes, metagenomes, and genomes at an unparalleled resolution, which can be essential in serving the provisioning ecosystem based on genetic material relevance to humans (Van Dijk et al., 2018). Plant science is taking advantage of the improvements in the field of DNA sequencing that are boosted by biomedical research intensives and influencing biodiversity protection and plant breeding (Delseny et al., 2010). The recent revolutions in genomics, combining precise phenotypes and high-throughput technology, facilitate the identification of agronomic trait-controlling genes. For example, crop wild relatives (CWRs) provide potential for the domestication of new individuals. The annotation and assembly of high-quality crop plants provide new research targets, such as cis-regulatory regions and long noncoding RNAs (lncRNAs). These advancements may also improve the selection for plant cultivation, and in the future, the crops can approach the demands of the changing climate and growing population (Pourkheirandish et al., 2020).

### 3.1. Genome Editing and Mutation Breeding

The leading causes of biodiversity degradation are habitat and species loss, invasive species, overexploitation, pollution, and climate change. Biodiversity conservation, along with innovation in plant breeding, has become a global concern requiring a comprehensive and integrated approach. There are different recently adopted modern techniques and strategies to develop novel plant types and conserve plant biodiversity (Ebert, 2020; Engels & Thormann, 2020). Numerous gene editing tools, such as zinc-finger nucleases (ZFNs) (Basak et al., 2021), transcription-activator-like effector nucleases (TALENs) (Becker & Boch, 2021) and clustered regulatory interspaced short palindromic repeat (CRISPR)/CRISPR-associated 9 protein (Cas9) systems (Bessoltane et al., 2022). Based on biological cutting techniques, they have a wide range of applications in crop breeding and improvement. ZFNs are created by fusing Fok I endonuclease (Osakabe et al., 2010), which has non-specific cutting activity, with a synthetically altered zinc-finger protein (ZFP), which has specific binding activity. A number of the amino acid residues on ZFP's helix directly contribute to the recognition of the target site and can pair with bases nearby (Novak, 2019). Because TALENs use the transcription activator-like effector (TALE) as their binding domain and the binding function mostly depends on highly changeable amino acids at the 12th and 13th positions on the TALE, they vary structurally from ZFPs. A single base can be specifically identified when combined with them (Barnett, 2018). One-guide RNA (sgRNA) and Cas9 were combined to create the CRISPR/Cas9 system. Today's extensive use of gene editing technology for crop enhancement is largely made possible by the development of the CRISPR/Cas9 system (Zhu et al., 2020). After undergoing particular alteration, the sgRNA can bind to the target location (Zhang et al., 2021). The Cas9 protein's HNH and RuvC domains can cleave DNA strands with protospacer-associated motif (PAM) sites, which are complementary to the sgRNA. The PAM location is close to the cutting site. Base editors (BEs) and prime editors (PEs) have been introduced, which has further enhanced the CRISPR/Cas system (Jacinto et al., 2020).

EvolvR system, which can continuously diversify all nucleotides within a tunable window length at user-defined loci in bacteria, has been developed based on nCas9-recruitment of error-prone DNA polymerases. It allows the identification of ribosomal mutations that confer resistance to the antibiotic spectinomycin in *E. coli* (Halperin et al., 2018). CRISPR-Cas13 knocks down RNAs using RNA-guided RNA-targeting CRISPR-Cas effector Cas13a. Direct adenosine-to-inosine deaminase activity performed by ADAR2 (adenosine deaminase acting on RNA type 2) can be targeted toward the transcripts produced by mammalian cells (Matsoukas, 2018). High-throughput mapping of genetic variants can be achieved using CRISPR-enabled trackable genome engineering (CREATE). CREATE cassette designer can be used to design PAM sequences near the target sites of interest. It can rapidly design >105 cassettes, which in turn increases editing efficiency (Ren et al., 2020).

RNA Interference Technology in maize silenced the *Se1* gene, which increased the soluble sugar content by

affecting the starch metabolism of the endosperm, making the mutant maize more favorable for intestinal digestion (Zhang et al., 2019). (Pérez et al., 2019) used *MIMI168* to inhibit the expression of microRNA168 in rice, increase the expression of the *AGO1* gene, and then increase the number of panicles. Compared with the control rice, the yield was increased by 30–40%. Knockout of the *OsAAP6* gene using the CRISPR/Cas9 system then promoted rice tillering, thereby increasing rice yield (Lu et al., 2018). Zhang et al. (2021) used the CRISPR/Cas9 system to edit the *Ppo* gene of wheat, which affected the expression of polyphenol oxidase, inhibited dough browning, and improved the quality of wheat. These initiatives have advanced the development of gene editing technology and given us a powerful weapon for accelerating the domestication of crops, enlarging the crop gene pool and enhancing crop output and quality.

### 3.2. Future Endeavors of Plant Breeding to Achieve Sustainable Crop Production with Maximum Biodiversity

Climate change is a global phenomenon and a threat to food security in the future. Plant breeders emphasize the higher yield in crop plants with a broader genetic base. To achieve goals in plant breeding, the use of genetic resources is very crucial. It makes it possible to gain maximum genetic diversity to boost crop yield with climate-resilient lines. The idea of climate-resilient crops is growing very fast in the modern era as it protects the future of food security. Along with a broader genetic base and higher yield in changing climates, breeders are also trying to develop nutrient-enriched crops. For this purpose, wheat, corn, and rice have been improved in the recent past to get nutrient-enriched grains. Flavr Savr and purple tomatoes are other examples of nutrient-enriched crops to fight against climate change. All these developments were only made through the utilization of different genetic resources. Scientists were not interested in conserving genetic resources after developing high-yielding crops, but now they are firstly conserving the genetic base of any breeding material for the future and then focusing on their desired products. So, the application of modern breeding techniques to develop highly efficient crops from genetic resources is the only way to fight future hunger.

## 4. CONCLUSION

Conserving and utilizing different genetic resources for crop improvement to cope effects of climate changes on agricultural production. Plant breeders and geneticists play a vital role in this work. Genetic resources are the source of a broad genetic base for any crop to be developed as climate resilient in the future. Modern plant breeding techniques make it more proficient to get higher yield and nutrient enriched crops by utilizing conserved genetic resources to reduce hunger in the scenario of climate change. Recently, different modern techniques have been developed to target a specific trait in a crop or accumulate the desired traits in a single line. In the future, these lines could serve as climate-resilient crops. Genetic resources are also being conserved using different conservation techniques. When a breeder gets germplasm of any crop from genetic resources later, they'll have to submit the equal germplasm after multiplication. So, the conservation of genetic resources and their utilization in modern breeding programs are the ghosts to making highly nutritious, high-yielding crops with a broader genetic base.

## DECLARATIONS

**Funding:** There is no funding for this review article.

**Acknowledgement:** We are thankful to Dr. Muhammad Imtiaz Rashid from King Abdullah University, Saudi Arabia, and Miss Maria Batool from Huazhong Agricultural University, China, for their assistance in improving the language and clarity of this review paper.

**Conflicts of Interest:** The authors have no relevant financial or non-financial interests to disclose.

**Data Availability:** The current review article is not available in any repository and has not been submitted before in any journal.

**Author's Contribution:** All authors contributed equally in designing outlines, collecting materials, writing manuscripts, and reviewing them.

**Generative AI Statement:** The authors declare that no Gen AI/DeepSeek was used in the writing/creation of this manuscript.

**Publisher's Note:** All claims stated in this article are exclusively those of the authors and do not necessarily represent those of their affiliated organizations or those of the publisher, the editors, and the reviewers. Any product that may

**Citation:** Manzoor Z, Shahzad U, Qadir MS, Jahan MS, Hassan Z, Parveen I, Nawaz MS, Raza A, Shahnaz Z, Bibi M, Aslam U, Khalid I, Nadeem S, Khan MY and Rasool MG, 2025. Utilizing crop genetic resources in modern breeding to achieve sustainable goals in a changing climate. *Agrobiological Records* 20: 74-83. <https://doi.org/10.47278/journal.abr/2025.021>

be evaluated/assessed in this article or claimed by its manufacturer is not guaranteed or endorsed by the publisher/editors.

## REFERENCES

- Abberton, M., Batley, J., Bentley, A., Bryant, J., Cai, H., Cockram, J., & De Pace, C. (2016). Global agricultural intensification during climate change: A role for genomics. *Plant Biotechnology Journal*, 14(4), 1095–1098. <https://doi.org/10.1111/pbi.12467>
- Alex Buerkle, C., & Gompert, Z. (2013). Population genomics based on low coverage sequencing: How low should we go? *Molecular Ecology*, 22(11), 3028–3035. <https://doi.org/10.1111/mec.12105>
- Andrews, K. R., Good, J. M., Miller, M. R., Luikart, G., & Hohenlohe, P. A. (2016). Harnessing the power of RADseq for ecological and evolutionary genomics. *Nature Reviews Genetics*, 17(2), 81–92. <https://doi.org/10.1038/nrg.2015.28>
- Arora, N. K. (2018). Biodiversity conservation for sustainable future. In Vol. 1 (pp. 109–111). Springer. [https://doi.org/10.1007/978-981-10-6896-0\\_16](https://doi.org/10.1007/978-981-10-6896-0_16)
- Awan, F. S., Sadia, B., Altaf, J., Habib, M., Hameed, K., & Hussain, S. (2021). Genetic variability through induced mutation. In *Genetic Variation*. IntechOpen. <https://doi.org/10.5772/intechopen.95027>
- Baidyussen, A., Jatayev, S., Khassanova, G., Amantayev, B., Sereda, G., Sereda, S., Gupta, N. K., Gupta, S., Schramm, C., Anderson, P., & Jenkins, C. L. (2021). Expression of specific alleles of zinc-finger transcription factors, HvSAP8 and HvSAP16, and corresponding SNP markers, are associated with drought tolerance in barley populations. *International Journal of Molecular Sciences*, 22(22), 12156. <https://doi.org/10.3390/ijms222212156>
- Bardak, A., & Bölek, Y. (2012). Genetic diversity of diploid and tetraploid cottons determined by SSR and ISSR markers. *Turkish Journal of Field Crops*, 17(2), 139–144.
- Barnett, P. (2018). Transcription activator-like effector nucleases (TALENs): A new, important, and versatile gene editing technique with a growing literature. *Science & Technology Libraries*, 37(1), 100–112. <https://doi.org/10.1080/0194262X.2018.1424114>
- Basak, N., Verma, V. C., Kumar, R., & Kumar, G. (2021). Mechanism of ZFN-mediated genome editing: Scope and opportunities. In *Genome editing in plants* (pp. 13–27). CRC Press.
- Becker, S., & Boch, J. (2021). TALE and TALEN genome editing technologies. *Gene and Genome Editing*, 2, 100007. <https://doi.org/10.1016/j.ggedit.2021.100007>
- Beissinger, M. R. (2022). *The Revolutionary City: Urbanization and the global transformation of rebellion*. Princeton University Press. <https://doi.org/10.1515/9780691224756>
- Bessoltane, N., Charlot, F., Guyon-Debast, A., Charif, D., Mara, K., Collonnier, C., & Nogu  , F. (2022). Genome-wide specificity of plant genome editing by both CRISPR–Cas9 and TALEN. *Scientific Reports*, 12(1), 13034. <https://doi.org/10.1038/s41598-022-13034-2>
- Brooks, C., Nekrasov, V., Lippman, Z. B., & Van Eck, J. (2014). Efficient gene editing in tomato in the first generation using the clustered regularly interspaced short palindromic repeats/CRISPR-associated9 system. *Plant Physiology*, 166(3), 1292–1297. <https://doi.org/10.1104/pp.114.247148>
- Bukhari, S. A. R., Saeed, M., & Briddon, R. W. (2021). Use of CRISPR/Cas system to create resistance to cotton diseases. In *Cotton Precision Breeding* (pp. 329–350). Cham: Springer International Publishing. [https://doi.org/10.1007/978-3-030-82524-4\\_17](https://doi.org/10.1007/978-3-030-82524-4_17)
- Cortes-Olmos, C., Vilanova, S., Pascual, L., Rosell  , J., & Cebolla-Cornejo, J. (2015). SNP markers applied to the characterization of Spanish tomato (*Solanum lycopersicum* L.) landraces. *Scientia Horticulturae*, 194, 100–110. <https://doi.org/10.1016/j.scienta.2015.07.036>
- Dahab, A. A., Saeed, M., Mohamed, B. B., Ashraf, M. A., Puspito, A. N., Shahid, K. S. B. A. A., & Husnain, T. (2013). Genetic diversity assessment of cotton (*Gossypium hirsutum* L.) genotypes from Pakistan using simple sequence repeat markers. *Australian Journal of Crop Science*, 7(2), 261–267.
- de Boef, W. S., Subedi, A., Peroni, N., Thijssen, M., & O’Keeffe, E. (2013). *Community biodiversity management: Promoting resilience and the conservation of plant genetic resources*. Routledge.
- De Groot, G. A., Nowak, C., Skrbin  ek, T., Andersen, L. W., Aspi, J., Fumagalli, L., & Liberg, O. (2016). Decades of population genetic research reveal the need for harmonization of molecular markers: The grey wolf *Canis lupus* as a case study. *Mammal Review*, 46(1), 44–59. <https://doi.org/10.1111/mam.12050>
- Delseny, M., Han, B., & Hsing, Y. I. (2010). High throughput DNA sequencing: The new sequencing revolution. *Plant Science*, 179(5), 407–422. <https://doi.org/10.1016/j.plantsci.2010.06.011>
- Dreisigacker, S., Zhang, P., Warburton, M., Skovmand, B., Hoisington, D., & Melchinger, A. (2005). Genetic diversity among and within CIMMYT wheat landrace accessions investigated with SSRs and implications for plant genetic resources management. *Crop Science*, 45(2), 653–661. <https://doi.org/10.2135/cropsci2005.0653>
- Du, H., Zeng, X., Zhao, M., Cui, X., Wang, Q., Yang, H., & Yu, D. (2016). Efficient targeted mutagenesis in soybean by TALENs and CRISPR/Cas9. *Journal of Biotechnology*, 217, 90–97. <https://doi.org/10.1016/j.jbiotec.2015.11.005>
- Duitama, J., Silva, A., Sanabria, Y., Cruz, D. F., Quintero, C., Ballen, C., & Torres, E. (2015). Whole genome sequencing of elite rice cultivars as a comprehensive information resource for marker-assisted selection. *PLOS ONE*, 10(4), e0124617. <https://doi.org/10.1371/journal.pone.0124617>
- Ebert, A. W. (2020). The role of vegetable genetic resources in nutrition security and vegetable breeding. *Plants*, 9(6), 736. <https://doi.org/10.3390/plants9060736>

**Citation:** Manzoor Z, Shahzad U, Qadir MS, Jahan MS, Hassan Z, Parveen I, Nawaz MS, Raza A, Shahnaz Z, Bibi M, Aslam U, Khalid I, Nadeem S, Khan MY and Rasool MG. 2025. Utilizing crop genetic resources in modern breeding to achieve sustainable goals in a changing climate. *Agrobiological Records* 20: 74-83. <https://doi.org/10.47278/journal.abr/2025.021>

- Ebert, A. W., & Engels, J. M. (2020). Plant biodiversity and genetic resources matter! *Plants*, 9, 1706. <https://doi.org/10.3390/plants9121706>
- Engels, J. M., & Thormann, I. (2020). Main challenges and actions needed to improve conservation and sustainable use of our crop wild relatives. *Plants*, 9(8), 968. <https://doi.org/10.3390/plants9080968>
- Fasoula, D. A., Ioannides, I. M., & Omirou, M. (2020). Phenotyping and plant breeding: Overcoming the barriers. *Frontiers in Plant Science*, 10, 1713. <https://doi.org/10.3389/fpls.2019.01713>
- Głowacka, K., Kromdijk, J., Kucera, K., Xie, J., Cavanagh, A. P., Leonelli, L., & Long, S. P. (2018). Photosystem II subunit S overexpression increases the efficiency of water use in a field-grown crop. *Nature Communications*, 9(1), 1–9. <https://doi.org/10.1038/s41467-018-03674-z>
- Gollin, D. (2020). Conserving genetic resources for agriculture: Economic implications of emerging science. *Food Security*, 12(5), 919–927. <https://doi.org/10.1007/s12571-020-01054-8>
- Hakeem, K. R. (2015). Crop production and global environmental issues. Springer. <https://doi.org/10.1007/978-3-319-23162-4>
- Halperin, S. O., Tou, C. J., Wong, E. B., Modavi, C., Schaffer, D. V., & Dueber, J. E. (2018). CRISPR-guided DNA polymerases enable diversification of all nucleotides in a tunable window. *Nature*, 560(7717), 248–252. <https://doi.org/10.1038/s41586-018-0384-8>
- Han, J., Guo, B., Guo, Y., Zhang, B., Wang, X., & Qiu, L.-J. (2019). Creation of early flowering germplasm of soybean by CRISPR/Cas9 technology. *Frontiers in Plant Science*, 10, 1446. <https://doi.org/10.3389/fpls.2019.01446>
- Haque, M. A., Rafii, M. Y., Yusoff, M. M., Ali, N. S., Yusuff, O., Datta, D. R., Anisuzzaman, M., & Ikbal, M. F. (2021). Recent advances in rice varietal development for durable resistance to biotic and abiotic stresses through marker-assisted gene pyramiding. *Sustainability*, 13(19), 10806. <https://doi.org/10.3390/su131910806>
- Hartati, R. S., Suhesti, S., Wulandari, S., Ardana, I. K., & Yunita, R. (2021). In-vitro selection of sugarcane (*Saccharum officinarum* L.) putative mutant for drought stress. In *IOP Conference Series: Earth and Environmental Science*, 653(1), 012135. <https://doi.org/10.1088/1755-1315/653/1/012135>
- IFPRI, 2009. IFPRI Annual Report 2008-2009. International Food Policy Research Institute. <https://ideas.repec.org/p/fpr/annrep/2009.html>
- Jacinto, F. V., Link, W., & Ferreira, B. I. (2020). CRISPR/Cas9-mediated genome editing: From basic research to translational medicine. *Journal of Cellular and Molecular Medicine*, 24(7), 3766–3778. <https://doi.org/10.1111/jcmm.15118>
- Jeeterwal, R. C., & Nehra, A. (2018). Principles of quality seed production of cereal crops. [Publisher not provided — please supply if known]
- Jena, K., & Mackill, D. (2008). Molecular markers and their use in marker-assisted selection in rice. *Crop Science*, 48(4), 1266–1276. <https://doi.org/10.2135/cropsci2007.05.0279>
- Jinek, M., Chylinski, K., Fonfara, I., Hauer, M., Doudna, J. A., & Charpentier, E. (2012). A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science*, 337(6096), 816–821. <https://doi.org/10.1126/science.1225829>
- Kasso, M., & Balakrishnan, M. (2013). Ex situ conservation of biodiversity with particular emphasis to Ethiopia. *International Scholarly Research Notices*, 2013, 985037. <https://doi.org/10.5402/2013/985037>
- Kim, C. G. (2012). The impact of climate change on the agricultural sector: Implications of the agro-industry for low carbon, green growth strategy and roadmap for the East Asian region. Economic and Social Commission for Asia and the Pacific (ESCAP): Bangkok, Thailand.
- Kim, T.-S., He, Q., Kim, K.-W., Yoon, M.-Y., Ra, W.-H., Li, F. P., & Choi, B. (2016). Genome-wide resequencing of KRICE\_CORE reveals their potential for future breeding, as well as functional and evolutionary studies in the post-genomic era. *BMC Genomics*, 17(1), 1–13. <https://doi.org/10.1186/s12864-016-2771-4>
- Kim, Y. X., Son, S., Lee, S., Jung, E., Lee, Y., Sung, J., & Lee, C. (2021). Combined effects of nutrients × water × light on metabolite composition in tomato fruits (*Solanum lycopersicum* L.). *Plants*, 10(7), 1437. <https://doi.org/10.3390/plants10071437>
- Kjeldsen, S. R., Zenger, K. R., Leigh, K., Ellis, W., Tobey, J., Phalen, D., & Raadsma, H. W. (2016). Genome-wide SNP loci reveal novel insights into koala (*Phascolarctos cinereus*) population variability across its range. *Conservation Genetics*, 17(2), 337–353. <https://doi.org/10.1007/s10592-015-0785-9>
- Koonin, E. V., Makarova, K. S., & Zhang, F. (2017). Diversity, classification and evolution of CRISPR-Cas systems. *Current Opinion in Microbiology*, 37, 67–78. <https://doi.org/10.1016/j.mib.2017.05.008>
- Kosgei, A. J., Kimurto, P. K., Gaur, P. M., Yeboah, M. A., Ofei, S. K., & Danquah, E. Y. (2022). Introgression of drought tolerance root traits into Kenyan commercial chickpea varieties using marker assisted backcrossing. *African Crop Science Journal*, 30(1), 31–50. <https://doi.org/10.4314/acsj.v30i1.3>
- Lei, Q.-Y., Zhou, J.-J., Xiong, Y., Zhang, W.-H., Luo, J., & Long, C.-L. (2021). Genetic diversity evaluation and conservation of Kam fragrant glutinous rice (*Oryza sativa* L.) germplasm in Southeast Guizhou, China. *Plants*, 10(9), 1898. <https://doi.org/10.3390/plants10091898>
- Li, H., Rasheed, A., Hickey, L. T., & He, Z. (2018). Fast-forwarding genetic gain. *Trends in Plant Science*, 23(3), 184–186. <https://doi.org/10.1016/j.tplants.2018.01.007>
- Lu, K., Wu, B., Wang, J., Zhu, W., Nie, H., Qian, J., & Fang, Z. (2018). Blocking amino acid transporter OsAAP3 improves grain yield by promoting outgrowth buds and increasing tiller number in rice. *Plant Biotechnology Journal*, 16(10), 1710–1722. <https://doi.org/10.1111/pbi.12907>
- Mashilo, J., Shimelis, H., & Odindo, A. (2017). Phenotypic and genotypic characterization of bottle gourd [*Lagenaria siceraria* (Molina) Standl.] and implications for breeding: A review. *Scientia Horticulturae*, 222, 136–144. <https://doi.org/10.1016/j.scienta.2017.04.006>

**Citation:** Manzoor Z, Shahzad U, Qadir MS, Jahan MS, Hassan Z, Parveen I, Nawaz MS, Raza A, Shahnaz Z, Bibi M, Aslam U, Khalid I, Nadeem S, Khan MY and Rasool MG. 2025. Utilizing crop genetic resources in modern breeding to achieve sustainable goals in a changing climate. *Agrobiological Records* 20: 74-83. <https://doi.org/10.47278/journal.abr/2025.021>



- Matsoukas, I. G. (2018). Commentary: RNA editing with CRISPR-Cas13. *Frontiers in Genetics*, 9, 134. <https://doi.org/10.3389/fgene.2018.00134>
- Mengist, W., Soromessa, T., & Feyisa, G. L. (2022). Estimating the total ecosystem services value of Eastern Afromontane biodiversity hotspots in response to landscape dynamics. *Environmental and Sustainability Indicators*, 14, 100178. <https://doi.org/10.1016/j.indic.2022.100178>
- Michel, S., Ametz, C., Gungor, H., Akgöl, B., Epure, D., Grausgruber, H., & Buerstmayr, H. (2017). Genomic assisted selection for enhancing line breeding: Merging genomic and phenotypic selection in winter wheat breeding programs with preliminary yield trials. *Theoretical and Applied Genetics*, 130(2), 363–376. <https://doi.org/10.1007/s00122-016-2820-5>
- Mohapatra, K. K., Singh, S. K., & Patra, A. (2021). Influence of varying levels of zinc on yield and zinc biofortification in hybrid rice (*Oryza sativa* L.) grown in moderate zinc soil. *Journal of the Indian Society of Soil Science*, 69(2), 220–223. <https://doi.org/10.5958/0974-0228.2021.00034.7>
- Mondal, S., & Palit, D. (2022). Challenges in natural resource management for ecological sustainability. In *Natural resources conservation and advances for sustainability* (pp. 29–59). Elsevier. <https://doi.org/10.1016/B978-0-323-85750-1.00002-4>
- Munda, S., Saikia, R. J., Begum, T., Bhandari, S., Gogoi, A., Sarma, N., & Lal, M. (2022). Evaluation of genetic diversity based on microsatellites and phytochemical markers of core collection of *Cymbopogon winterianus* Jowitt germplasm. *Plants*, 11(4), 528. <https://doi.org/10.3390/plants11040528>
- Nelson, E., Mendoza, G., Regetz, J., Polasky, S., Tallis, H., Cameron, D., & Kareiva, P. M. (2009). Modeling multiple ecosystem services, biodiversity conservation, commodity production, and tradeoffs at landscape scales. *Frontiers in Ecology and the Environment*, 7(1), 4–11. <https://doi.org/10.1890/080023>
- Novak, S. (2019). Plant biotechnology applications of zinc finger technology. In *Transgenic plants* (pp. 295–310). Springer. [https://doi.org/10.1007/978-3-030-04904-0\\_15](https://doi.org/10.1007/978-3-030-04904-0_15)
- Nyadanu, D., Adu Amoah, R., Obeng, B., Kwarteng, A., Akromah, R., Aboagye, L., & Adu-Dapaah, H. (2017). Ethnobotany and analysis of food components of African locust bean (*Parkia biglobosa* (Jacq.) Benth.) in the transitional zone of Ghana: Implications for domestication, conservation, and breeding of improved varieties. *Genetic Resources and Crop Evolution*, 64(6), 1231–1240. <https://doi.org/10.1007/s10722-016-0435-1>
- Ortiz-Bobea, A., Ault, T. R., Carrillo, C. M., Chambers, R. G., & Lobell, D. B. (2021). Anthropogenic climate change has slowed global agricultural productivity growth. *Nature Climate Change*, 11(4), 306–312. <https://doi.org/10.1038/s41558-021-01000-1>
- Osakabe, K., Osakabe, Y., & Toki, S. (2010). Site-directed mutagenesis in *Arabidopsis* using custom-designed zinc finger nucleases. *Proceedings of the National Academy of Sciences*, 107(26), 12034–12039. <https://doi.org/10.1073/pnas.1000235107>
- Panda, D., Barik, J., & Sarkar, R. K. (2021). Recent advances of genetic resources, genes, and genetic approaches for flooding tolerance in rice. *Current Genomics*, 22(1), 41–58. <https://doi.org/10.2174/1389202921999201228150023>
- Pastorino, M. J., & Marchelli, P. (2022). Genetic resources: The base material for managing nature. In: *Imperiled: The Encyclopedia of Conservation*. Elsevier, pp: 77-87. <https://doi.org/10.1016/B978-0-12-821139-7.00192-6>
- Paulino, J. F. C., de Almeida, C. P., Bueno, C. J., Song, Q., Fritsche-Neto, R., Carbonell, S. A. M., Chiorato, A. F., & Benchimol-Reis, L. L. (2021). Genome-wide association study reveals genomic regions associated with fusarium wilt resistance in common bean. *Genes*, 12(5), 765. <https://doi.org/10.3390/genes12050765>
- Pérez, L., Soto, E., Farré, G., Juanos, J., Villorquina, G., Bassie, L., & Rojas, J. A. (2019). CRISPR/Cas9 mutations in the rice Waxy/GBSSI gene induce allele-specific and zygoty-dependent feedback effects on endosperm starch biosynthesis. *Plant Cell Reports*, 38(3), 417–433. <https://doi.org/10.1007/s00299-019-02380-0>
- Pourkheirandish, M., Golicz, A. A., Bhalla, P. L., & Singh, M. B. (2020). Global role of crop genomics in the face of climate change. *Frontiers in Plant Science*, 11, 922. <https://doi.org/10.3389/fpls.2020.00922>
- Rahman, M. M., Mahmud, M. A. A., & Shahidullah, M. (2017). Socioeconomics of biodiversity conservation in the protected areas: A case study in Bangladesh. *International Journal of Sustainable Development & World Ecology*, 24(1), 65–72. <https://doi.org/10.1080/13504509.2016.1169453>
- Rana, N., Rahim, M. S., Kaur, G., Bansal, R., Kumawat, S., Roy, J., Deshmukh, R., Sonah, H., & Sharma, T. R. (2020). Applications and challenges for efficient exploration of omics interventions for the enhancement of nutritional quality in rice (*Oryza sativa* L.). *Critical Reviews in Food Science and Nutrition*, 60(19), 3304–3320. <https://doi.org/10.1080/10408398.2019.1685454>
- Ray, D. K., Mueller, N. D., West, P. C., & Foley, J. A. (2013). Yield trends are insufficient to double global crop production by 2050. *PLOS ONE*, 8(6), e66428. <https://doi.org/10.1371/journal.pone.0066428>
- Razzaq, A., Kaur, P., Akhter, N., Wani, S. H., & Saleem, F. (2021). Next-generation breeding strategies for climate-ready crops. *Frontiers in Plant Science*, 12, 637366. <https://doi.org/10.3389/fpls.2021.637366>
- Reardon, S. (2016). Welcome to the CRISPR zoo. *Nature News*, 531(7593), 160–163. <https://doi.org/10.1038/531160a>
- Reddy, B. J., Mandal, R., Chakroborty, M., Hijam, L., & Dutta, P. (2018). A Review on Potato (*Solanum Tuberosum* L.) and its Genetic Diversity. *International Journal of Genetics*, 10(2), 360. <https://doi.org/10.9735/0975-2862.10.2.360-364>
- Ren, J., Lee, J., & Na, D. (2020). Recent advances in genetic engineering tools based on synthetic biology. *Journal of Microbiology*, 58(1), 1–10. <https://doi.org/10.1007/s12275-020-9381-4>
- Sánchez-León, S., Gil-Humanes, J., Ozuna, C. V., Giménez, M. J., Sousa, C., Voytas, D. F., & Barro, F. (2018). Low-gluten, nontransgenic wheat engineered with CRISPR/Cas9. *Plant Biotechnology Journal*, 16(4), 902–910. <https://doi.org/10.1111/pbi.12837>

- Sharmin, A., Hoque, M. E., Haque, M. M., & Khatun, F. (2018). Molecular diversity analysis of some chilli (*Capsicum* spp.) genotypes using SSR markers. *American Journal of Plant Sciences*, 9(3), 368–378. <https://doi.org/10.4236/ajps.2018.93029>
- Sim, S. C., Van Deynze, A., Stoffel, K., Douches, D. S., Zarka, D., Ganai, M. W., ... Francis, D. M. (2012). High-density SNP genotyping of tomato (*Solanum lycopersicum* L.) reveals patterns of genetic variation due to breeding. *PLOS ONE*, 7(9), e45520. <https://doi.org/10.1371/journal.pone.0045520>
- Singh, M., Nara, U., Kumar, A., Thapa, S., Jaswal, C., & Singh, H. (2021). Enhancing genetic gains through marker-assisted recurrent selection: From phenotyping to genotyping. *Cereal Research Communications*, 1–16. <https://doi.org/10.1007/s42976-021-00142-y>
- Sun, Q., Zhao, Y., Zhang, Y., Chen, S., Ying, Q., Lv, Z., Che, X., & Wang, D. (2022). Heat stress may cause a significant reduction of rice yield in China under future climate scenarios. *Science of the Total Environment*, 818, 151746. <https://doi.org/10.1016/j.scitotenv.2021.151746>
- Svitashev, S., Young, J. K., Schwartz, C., Gao, H., Falco, S. C., & Cigan, A. M. (2015). Targeted mutagenesis, precise gene editing, and site-specific gene insertion in maize using Cas9 and guide RNA. *Plant Physiology*, 169(2), 931–945. <https://doi.org/10.1104/pp.15.00793>
- Todd, E. V., Black, M. A., & Gemmell, N. J. (2016). The power and promise of RNA-seq in ecology and evolution. *Molecular Ecology*, 25(6), 1224–1241. <https://doi.org/10.1111/mec.13526>
- Torchio, M. F., Lucia, U., & Grisolia, G. (2020). Economic and human features for energy and environmental indicators: A tool to assess countries' progress towards sustainability. *Sustainability*, 12(22), 9716. <https://doi.org/10.3390/su12229716>
- Van Dijk, E. L., Jaszczyszyn, Y., Naquin, D., & Thermes, C. (2018). The third revolution in sequencing technology. *Trends in Genetics*, 34(9), 666–681. <https://doi.org/10.1016/j.tig.2018.05.008>
- Verhoeven, K. J., Vonholdt, B. M., & Sork, V. L. (2016). Epigenetics in ecology and evolution: What we know and what we need to know. *Molecular Ecology*, 25(8), 1631–1638. <https://doi.org/10.1111/mec.13617>
- Verma, A., & Singh, G. P. (2020). Combining AMMI and mean yield of wheat genotypes evaluated under rainfed conditions of northern hills zone for stability analysis. *International Journal of Bioresource and Stress Management*, 11(6), 590–600. <https://doi.org/10.23910/I.2020.2162b>
- Wang, W., & Li, J. (2021). In-situ conservation of biodiversity in China: Advances and prospects. *Biodiversity Science*, 29(2), 133. <https://doi.org/10.17520/biods.2021085>
- Weeks, D. P., Spalding, M. H., & Yang, B. (2016). Use of designer nucleases for targeted gene and genome editing in plants. *Plant Biotechnology Journal*, 14(2), 483–495. <https://doi.org/10.1111/pbi.12448>
- Wellband, K., Baillie, S. M., Bentzen, P., & Bernatchez, L. (2021). Genetic diversity. In *The Lake Charr *Salvelinus namaycush*: Biology, ecology, distribution, and management* (pp. 119–165). Springer. [https://doi.org/10.1007/978-3-030-62203-8\\_6](https://doi.org/10.1007/978-3-030-62203-8_6)
- Yugi, K., Kubota, H., Hatano, A., & Kuroda, S. (2016). Trans-omics: How to reconstruct biochemical networks across multiple 'omic' layers. *Trends in Biotechnology*, 34(4), 276–290. <https://doi.org/10.1016/j.tibtech.2015.12.013>
- Zafar, M. M., Ijaz, A., Anwar, Z., Iqbal, M. S., Zafar, S., Subhan, M., Seleiman, M. F., Alhammad, B. A., Aljabri, M., Qiao, F., & Shoukat, A. (2025). Harnessing genetic diversity in cotton for enhanced resilience against salt stress by using agro-physiological characters. *Plant Production Science*, 28(1), 51–68. <https://doi.org/10.1080/1343943X.2024.2439874>
- Zhang, N., Liao, Z., Wu, S., Nobis, M. P., Wang, J., & Wu, N. (2022). Impact of climate change on wheat security through an alternate host of stripe rust. *Food and Energy Security*, 11(1), e356. <https://doi.org/10.1002/fes3.356>
- Zhang, X., Haro von Mogel, K. J., Lor, V. S., Hirsch, C. N., De Vries, B., Kaeppler, H. F., Tracy, W. F., & Kaeppler, S. M. (2019). Maize sugary enhancer1 (*se1*) is a gene affecting endosperm starch metabolism. *Proceedings of the National Academy of Sciences*, 116(41), 20776–20785. <https://doi.org/10.1073/pnas.1902747116>
- Zhang, X., Xu, G., Cheng, C., Lei, L., Sun, J., Xu, Y., Deng, C., Dai, Z., Yang, Z., Chen, X., & Liu, C. (2021). Establishment of an *Agrobacterium*-mediated genetic transformation and CRISPR/Cas9-mediated targeted mutagenesis in hemp (*Cannabis sativa* L.). *Plant Biotechnology Journal*, 19(10), 1979–1987. <https://doi.org/10.1111/pbi.13611>
- Zhao, W., Lee, G. A., Kwon, S. W., Ma, K. H., Lee, M. C., & Park, Y. J. (2012). Development and use of novel SSR markers for molecular genetic diversity in Italian millet (*Setaria italica* L.). *Genes & Genomics*, 34, 51–57. <https://doi.org/10.1007/s13258-011-0197-z>
- Zhao, X., Meng, Z., Wang, Y., Chen, W., Sun, C., Cui, B., Cui, J., Yu, M., Zeng, Z., Guo, S., & Luo, D. (2017). Pollen magnetofection for genetic modification with magnetic nanoparticles as gene carriers. *Nature Plants*, 3(12), 956–964. <https://doi.org/10.1038/s41477-017-0063-z>
- Zhou, J., Peng, Z., Long, J., Sosso, D., Liu, B., Eom, J. S., Huang, S., Liu, S., Vera Cruz, C., Frommer, W. B., White, F. F., & Yang, B. (2015). Gene targeting by the TAL effector PthXo2 reveals cryptic resistance gene for bacterial blight of rice. *The Plant Journal : For Cell and Molecular Biology*, 82(4), 632–643. <https://doi.org/10.1111/tpj.12838>
- Zhou, Q., Zhou, C., Zheng, W., Mason, A. S., Fan, S., Wu, C., Fu, D., & Huang, Y. (2017). Genome-wide SNP markers based on SLAF-seq uncover breeding traces in rapeseed (*Brassica napus* L.). *Frontiers in Plant Science*, 8, 648. <https://doi.org/10.3389/fpls.2017.00648>
- Zhu, H., Li, C., & Gao, C. (2020). Applications of CRISPR–Cas in agriculture and plant biotechnology. *Nature Reviews Molecular Cell Biology*, 21(11), 661–677. <https://doi.org/10.1038/s41580-020-00288-x>