








USE OF MEDICINAL PLANTS AS ALTERNATIVE FOR THE CONTROL OF INTESTINAL PARASITOSIS: ASSESSMENT AND PERSPECTIVES

Lissette H. Degla ^{1,5}, Julienne Kuiseu ¹, Pascal A. Olounlade ^{1,3*}, Sabbas Attindehou ², Sylvie M. Hounzangbe-Adote ³, Patrick A. Eдорh ⁴ and Latifou Lagnika ⁵

¹Zootechnical Research and Livestock System Unit, Laboratory of Animal and Fisheries Sciences, Doctoral School of Agricultural and Water Sciences, National University of Agriculture, 01 BP: 55 Porto-Novo, Benin

²Animal Health and Biosecurity Research Unit, Laboratory of Animal and Fisheries Sciences, Doctoral School of Agricultural and Water Sciences, National University of Agriculture, 01 BP: 55 Porto-Novo, Benin

³Laboratory of Ethnopharmacology and Animal Health, Faculty of Agronomic Sciences, University of Abomey-Calavi, 01 BP 526 Cotonou, Benin

⁴Laboratory of Toxicology and Environmental Health (LATSE), University of Abomey-Calavi (UAC), 01 BP 526 Cotonou, Republic of Benin

⁵Laboratory of Biochemistry and Bioactive Natural Substances, Faculty of Science and Technology, University of Abomey-Calavi, 04BP 0320, Cotonou, Benin

*Corresponding author: abioudouno@yahoo.fr; polounlade@gouv.bj

ABSTRACT

Intestinal parasitosis is a significant health problem for animal husbandry, leading to economic losses. In the treatment of helminths, synthetic anthelmintics have long been used. However, cases of parasitic resistance to these anthelmintics have been reported worldwide. Solutions have been proposed to solve this problem. Among them is the use of medicinal plants with anthelmintic properties. This work suggests synthesizing the work carried out on evaluating the anthelmintic properties of medicinal plants used to treat intestinal parasitosis of small ruminants. According to the results of the ethnobotanical surveys reported, several medicinal plants are used by the populations to treat intestinal parasitosis of small ruminants. Evaluations of the anthelmintic properties in vitro and or in vivo of some of them have confirmed their potential to be used as an alternative for controlling intestinal parasitosis. However, these results obtained depend on the organ of the species used, the type of extract, and the application dose. Tannins, flavonoids, and essential oils are the secondary metabolites responsible for the anthelmintic activity of these medicinal plants with anthelmintic potential. The efficacy of a plant extract (or powder) also depends on the parasite used in the tests. The results of previous studies confirm the use of medicinal plants in the fight against intestinal parasitosis.

Keywords: Anthelmintic, Intestinal parasitosis, Medicinal plants, Resistance

Article History (2021-0876) || Received: 08 Aug 2021 || Revised: 01 Sep 2021 || Accepted: 05 Sep 2021 || Published Online: 13 Sep 2021

1. INTRODUCTION

Gastrointestinal parasitosis remains a real health problem for small ruminant livestock, especially because of the resistance developed by parasites against synthetic anthelmintics (Akoto et al. 2019; Nath et al. 2019; Rehman et al. 2021). Indeed, this state of affairs compromises the effectiveness of these drugs. Small ruminant breeding is an income-generating activity and therefore contributes to the national economy in developing countries like Benin. In pastures, small ruminants are more vulnerable to strongyles *Oesophagostoma venulosum*, *Chabertia ovina* (superfamily Strongyloidea), *Cooperia curticei*, *Nematodius battus*, *Trichostrongylus axei*, *Trichostrongylus colubriformis* and *Haemonchus contortus* (superfamily Trichostrongyloidea). The species of the superfamily Trichostrongyloidea are distinguished by their absent or rudimentary oral capsule and a highly developed copulatory bursa and are the most pathogenic strongyles (Aguerre 2019). Infections by these gastrointestinal parasites, lead to weight loss of animals, poor wool quality, generating economic losses and can compromise food security (Alowanou et al. 2015). It is then important to use other solutions to control gastrointestinal parasites. Among the proposed populations is the use of medicinal plants with anthelmintic potential. The knowledge about use of medicinal plants is a tradition that is perpetuated from one generation to another. Ecological zone, level of education and ethnicity are factors related to the knowledge and use of medicinal plants by herders in Benin (Dassou et al. 2015). As a result of exchanges with the populations, medicinal plants have been listed and some

have already been studied to evaluate their effectiveness on gastrointestinal parasites. However, the following question must be asked. What are or will be the challenges related to the use of medicinal plants in the treatment of gastrointestinal parasitosis? The research will have found its outcome when the breeder will have found relief or satisfaction through an effective treatment of gastrointestinal parasitosis of these animals at lower cost. It can be translated by the indication of effective plants and or the provision of improved traditional medicines accessible and at lower cost. It is with regard to all these observations, that the present study proposes to make the synthesis of the works in order to bring out the limits for a better use of the medicinal plants endowed with anthelmintic properties.

1.2. Parasite resistance to synthetic anthelmintics

Parasite resistance is defined as: the ability of a parasite population to resist concentrations of antiparasitic drugs that are usually lethal to individuals of that species. It is a mutational preadaptation whose determinism is genetic and inherited (Okombe and Pongombo 2013). Isotype 1 of the beta-tubulin gene SNPs is thought to be linked to helminth resistance to benzimidazole (Furtado et al. 2019). Factors favoring parasite resistance to anthelmintics are: frequency of treatments per time interval, adherence to dosage (under-dosing), exclusive use of a single anthelmintic or family of anthelmintics, grazing management (small ruminants). Other so called intrinsic factors are related to the life cycle of the parasites (short cycle), their prolificity (high) and pathogenicity (highly pathogenic) (Aguerre 2019). *Haemonchus contortus* parasite of small ruminants has a life cycle of 20 days and the female has the capacity to lay 5,000 to 10,000 eggs per day. It lodges in the abomasum and is hematophagous (Emery et al. 2016). These characteristics justify its resistance to synthetic anthelmintics. Resistance of other helminths to synthetic anthelmintics has also been reported. The Fig. 1 shows the three main lines of thought and alternative solutions in solving the difficulties related to the resistance of parasites to synthetic anthelmintics in the management of gastrointestinal parasitosis of small ruminants.




Fig. 1: Alternative solutions for the control of gastrointestinal parasitosis in small ruminants.

The interest of drying up the source of contamination is to block the biological cycle of the parasites in order to control the infestation of the pasture and reduce new infestations. It is done through prevention, i.e. healthy animals on clean pastures. In case of infections by gastrointestinal parasites, the treated animals will be transferred from contaminated pastures to clean pastures: this is the escape and finally the dilution. Another solution is to improve the host's resistance to parasites. This can be done by selecting NGIs resistant animals, by vaccination and or by improving the host's feed ration. The third alternative is to eliminate parasites within the host through the use of medicinal plants with anthelmintic properties (Kuisseu et al. 2021).

The use of medicinal plants with anthelmintic potentiality is the most promising alternative. The exploitation of medicinal plants will limit the presence of residues in the environment or in consumer products and would be more accessible to the population.

1.3. Medicinal plants with anthelmintic potential

Several medicinal plants are used by the populations in the treatment of gastrointestinal parasitosis of small ruminants (Table 1). These results are obtained on the basis of the endogenous knowledge of the populations. The knowledge and use of medicinal plants are a kind of tradition that is transmitted from one generation to another. Some of these plants with anthelmintic potential have been studied for their activities *in vitro* and / or *in vivo*. It is evaluated through tests on the different life stages (eggs, larvae and adult worms) of these parasites. *Haemonchus*

contortus is the most studied nematode of small ruminants and is justified by its veterinary importance. It is one of the most pathogenic parasites infecting small ruminants. In Benin, its prevalence is 55.56%. (Attindehou et al. 2012). It is a hematophagous nematode that lodges in the abomasum of animals and 2000 females can puncture 30 ml of blood per day from a sheep (Coyne and Smith 1992). Also, cases of resistance of the *Haemonchus contortus* parasite to synthetic anthelmintics have been reported.

Table 1: List of some medicinal plants traditionally used in the treatment of gastrointestinal parasitosis and tested in small ruminants

Family	Species	Parasite studied	In vitro	In vivo	Comments	References
Annonaceae	<i>Annona senegalensis</i>	<i>H. contortus</i>	-	+	The leaves of <i>A. senegalensis</i> appear to be more effective than other parts of the plant. The parasite spectrum sensitive to the different extracts of <i>A. senegalensis</i> is composed of strongyles (<i>H. contortus</i> , <i>Trychostrongylus</i> sp ...) and probably coccidia	Nguessan et al. (2017)
Anacardiaceae	<i>Spondias mombin</i>	<i>H. contortus</i>	+	+	Ethanollic and methanolic extracts of <i>S. mombin</i> leaves inhibited the migration of L3 larvae and the motility of adult <i>H. contortus</i> worms. However, the adult worms were significantly more sensitive after 6 hours of exposure to ethanolic extracts than methanolic extracts. <i>In vivo</i> , <i>S. mombin</i> leaf powder reduced the rate of egg excretion in parasitized animals.	Akouedegni et al. (2019)
Bignoniaceae	<i>Newbouldia laevis</i>	<i>T. colubriformis</i>	+	+	The tannins contained in the leaves of <i>N. laevis</i> are responsible for inhibiting the migration of L3 larvae. After digestion in the rumen the leaves of <i>N. laevis</i> would retain their anthelmintic activities	Olounladé et al. (2017); Sidi et al. (2017b)
Caesalpinioideae	<i>Cassia alata</i>	<i>H. contortus</i>	+	-	The aqueous and hydroacetic extract of <i>C. alata</i> leaves have ovicidal and larvicidal activities. The leaves are rich in steroidal and triterpenic compounds, anthracenosides, saponosides, polyphenols (tannins)	Tianhou et al. (2020)
	<i>Cassia italica</i>	<i>H. contortus</i>	+	-	The aqueous and ethanolic extracts of the leaves of <i>S. italica</i> contain polyphenols (flavonoid tannins). Aqueous and ethanolic extract of <i>S. italica</i> exhibited evidence of <i>in vitro</i> anthelmintic activity on eggs; infective larvae L3 and against adults of <i>H. contortus</i> .	Yongwa et al. (2020)
Chenopodiaceae	<i>Chenopodium ambrosioides</i>	<i>H. contortus</i> ; <i>O. columbianum</i>	+	-	The leaf powder of <i>C. ambrosioides</i> is rich in tannins, flavonoids, alkaloids, saponosides. aqueous extract of the leaves of <i>C. ambrosioides</i> has stronglycidal effects against <i>H. contortus</i> and <i>O. columbianum</i> .	Maiga et al. (2020)
Combretaceae	<i>Combretum glutinosum</i>	<i>H. contortus</i>	+	+	The leaves of <i>C. glutinosum</i> are rich in tannins, saponins flavonoids, alkaloids, glycosides, anthraquinones terpenes, reducing compounds. Acetonic and methanolic extracts of <i>C. glutinosum</i> leaves inhibited egg hatching, larval migration and adult worm motility.	Alowanou et al. (2019b)
Euphorbiaceae	<i>Bridelia ferruginea</i>	<i>H. contortus</i>	+	+	The leaves of <i>B. ferruginea</i> are rich in tannins, saponins, flavonoids, alkaloids, glycosides, anthraquinones, terpenes, reducing compounds. Acetonic and methanolic extracts of <i>B. ferruginea</i> leaves inhibited egg hatching, larval migration and adult worm motility. Acetonic extract and tannins from <i>B. ferruginea</i> leaves inhibited the emergence of <i>H. contortus</i> L3 larvae	Alowanou et al. (2019a); Tchetan et al. (2020)
Fabaceae	<i>Pterocarpus erinaceus</i>	<i>H. contortus</i>	+	-	The leaves of <i>P. erinaceus</i> are rich in anthracene derivatives, tannins, flavonoids, saponosides. Acetonic and methanolic extracts inhibited the migration of L3 larvae and the motility of adult <i>H. contortus</i> worms	Dedehou et al. (2014)

Lythraceae	<i>Punica granatum</i>	<i>H. contortus</i>	+	-	The roots of <i>P. granatum</i> contains Alkaloid, Saponin, Tannins, Flavonoids, Glycosides and Phenol. Methanolic extract of <i>P. granatum</i> inhibited egg hatch and motility of adult <i>H. contortus</i> worms	Ahmed et al. (2020)
Mimosoideae	<i>Parkia biglobosa</i>	<i>H. contortus</i>	+	-	Fruit pods are rich in alkaloids and tannins. The acetone extract of <i>P. biglobosa</i> fruit pods was more reactive on the migration of L3 larvae than the methanolic extract	Dedehou et al. (2014)
	<i>Albizia adianthifolia</i>	<i>H. contortus</i>	+	-	The hydroalcoholic extract of <i>A. adianthifolia</i> showed larvicidal activity on L3 larvae of <i>Haemonchus contortus</i>	Mesmin et al. (2018)
Moraceae	<i>Ficus exasperata</i>	<i>H. contortus</i>	+	-	Aqueous extract of <i>F. exasperata</i> inhibited adult worms motility	Houngnimassou n et al. (2017)
	<i>Ficus lutea</i>	<i>H. contortus</i>	+	-	The hydroalcoholic extract of <i>F. lutea</i> showed larvicidal activity on L3 larvae of <i>H. contortus</i>	Mesmin et al. (2018)
	<i>Morus Mesozygia</i>	<i>H. contortus</i>	+	-	The hydroalcoholic extract of <i>A. adianthifolia</i> showed ovicidal activity on the larval eggs of <i>H. contortus</i>	Mesmin et al. (2018)
Rubiaceae	<i>Mitragyna inermis</i>	<i>H. contortus</i> ; <i>T. colubriformis</i>	+	+	The leaves of <i>M. inermis</i> are rich in tannins, saponins, flavonoids, alkaloids, glycosides, anthraquinones, terpenes, reducing compounds. Aqueous extract of <i>M. inermis</i> leaves inhibited the motility of adult <i>T. colubriformis</i> worms. Acetonic and methanolic extracts of <i>M. inermis</i> leaves inhibited egg hatching, larval migration and adult worm motility. Acetonic extract and tannins from the leaves of <i>M. inermis</i> inhibited the emergence of L3 larvae of <i>H. contortus</i>	Alowanou et al. (2019a, b); Tchetan et al. (2020); Toklo et al. (2021)
Rutaceae	<i>Zanthoxylum zanthoxyloides</i>	<i>H. contortus</i> ; <i>T. Colubriformis</i>	+	+	The tannins contained in the leaves of <i>Z. zanthoxyloides</i> are responsible for inhibiting the migration of L3 larvae. After digestion in the rumen the leaves of <i>N. laevis</i> would retain their anthelmintic activities	Olounladé et al. (2017); Azando et al. (2017 Sidi et al. (2017b)
Combretaceae	<i>Terminalia Catappa</i>	<i>H. contortus</i>	+	-	Dichloromethane extract of <i>T. catappa</i> leaves inhibited the hatching of <i>H. contortus</i> eggs more than methanolic extract	Olukotun et al. (2018)
Asteraceae	<i>Artemisia herba-alba</i>	<i>H. contortus</i>	+	-	The flowers and aerial part of <i>A. herba-alba</i> contains alkaloid, saponin, tannins, flavonoids, glycosides and phenol. The methanolic extract of <i>A. herba-alba</i> inhibited egg hatch and motility of adult <i>H. contortus</i> worms	Ahmed et al. (2020)
Zingiberaceae	<i>Curcuma longa</i>	<i>H. contortus</i>	+	-	The hydroalcoholic extract of <i>C. longa</i> reduced the rate of egg excretion from infected animals	Nath et al. (2019)
Celastraceae	<i>Maytenus senegalensis</i>	<i>H. contortus</i>	+	-	Tannins are responsible for the anthelmintic activities of the aqueous extract of <i>M. senegalensis</i>	Zangueu et al. (2018)
Lamiaceae	<i>Thymus vulgaris</i> L.	<i>H. contortus</i>	+	+	Thymol, the main compound in the essential oil of <i>Thymus vulgaris</i> L, is believed to be responsible for its anthelmintic activity.	Ferreira et al. (2016)

Egg hatch tests, inhibition of larval migration, and adult worm motility are all techniques used to evaluate the efficacy of extracts on gastrointestinal nematodes. The egg hatch test is a test that evaluates the ability of an extract to act on egg hatching. The inhibition of the survival of adult worms induced by an extract is an important factor in the control of these parasites. Indeed, the disruption of their motility would result in difficulties in feeding (thus less blood depletion for hematophagous parasites) and mating (thus less egg excretion) (Zinsou 2015; Sobhy et al. 2021). The decrease in larval migration, could disrupt their settlement in the mucosal wall of the digestive tract (Dedehou et al. 2014; Tchetan et al. 2020). The tests performed *in vitro* allow a rapid evaluation of the anthelmintic properties of the different plants tested and the analysis of the mechanisms of action of these plants on the parasites (Akouedegni et al. 2019). *Z. zanthoxyloides*, *N. laevis*, *S. monbin*, *C. italitica*, *B. ferruginea*, *A. herba-alba*, *C. longa* are some of the species that demonstrated their anthelmintic activities *in vitro* through egg hatching, larval migration inhibition and adult worm motility tests. Within the organism, several other components such as the

metabolism of the medicinal plant (or its extract), the availability and concentration of the active principle(s) can influence the activity of the medicinal plant (or its extract), hence the importance of *in vivo* tests. Also, the doses and frequencies of ingestion of anthelmintic plants are important factors to control for good efficacy, as the concentrations in the digestate can vary from one portion of the digestive tract to another (Azando et al. 2011). Thus, results obtained during *in vitro* testing may not match the *in vivo* test (Akouedegni et al. 2019). *In vivo* tests performed on artificially infested sheep have demonstrated the therapeutic effect of extracts of *Z. zanthoxyloides*, *N. laevis*, *S. monbin*, *M. inermis* *A. senegalensis* (Azando et al. 2011; Azando et al. 2017; Akouedegni et al. 2019). *S. monbin* leaf powder and *Annona senegalensis* leaf extract administered to naturally infested sheep have been shown to be effective in reducing egg shedding with selectivity (Ademola et al. 2005; Nguessan et al. 2017). That is, the extracts are more active on some parasites than others. The aqueous extract of *Annona senegalensis* would not be effective on *Taenia* spp. (Nguessan et al. 2017). *Z. zanthoxyloides* and *N. laevis* powders would be more effective on *H. contortus* eggs than on *T. colubriformis* and *O. columbianum* eggs (Azando et al. 2011).

Table 2: List of some medicinal plants traditionally used in the treatment of gastrointestinal parasitosis and not yet tested in small ruminants

Family	Species	References
Acanthaceae	<i>Nelsonia canescens</i> (Lam.)	Dassou et al. (2015); Ogni et al. (2014)
Alliaceae	<i>Allium sativum</i>	Dassou et al. (2012); Ogni et al. (2014)
Amaranthaceae	<i>Amaranthus spinosus</i>	Ogni et al. (2014)
Anacardiaceae	<i>Anacardium occidentale</i>	Ogni et al. (2014)
	<i>Lansea acida</i>	Garba et al. (2019)
	<i>Mangifera indica</i>	Dassou et al. (2012); Ogni et al. (2014)
	<i>Sclerocarya bierrea</i>	Garba et al. (2019)
Apocynaceae	<i>Pleiocarpa pycnantha</i>	Dassou et al. (2012); Ogni et al. (2014)
Asclepiadaceae	<i>Leptadania hastata</i>	Garba et al. (2019)
Bignoniaceae	<i>Kigelia Africana</i>	Garba et al. (2019)
Bombacaceae	<i>Adansonia digitata</i>	Dassou et al. (2012); Ogni et al. (2014); Garba et al. (2019)
	<i>Ceiba pentandra</i>	Kabore et al. (2007)
Cachlospermaceae	<i>Cochlospermum planchonii</i>	Dassou et al. (2012); Ogni et al. (2014); Garba et al. (2019)
	<i>Cassia sieberiana</i>	Dassou et al. (2012); Garba et al. (2019)
Caesalpinioideae	<i>Tamarindus indica</i>	Ogni et al. (2014)
	<i>Boscia salicifolia</i>	Garba et al. (2019)
Capparaceae	<i>Boscia senegalensis</i>	Garba et al. (2019)
	<i>Cadaba farinosa</i>	Garba et al. (2019)
	<i>Crateva adansonii</i>	Dassou et al. (2012); Ogni et al. (2014)
	<i>Maerua crassifolia</i>	Garba et al. (2019)
	<i>Carica papaya</i>	Dassou et al. (2012); Attindehou et al. (2012); Sidi et al. (2017a); Islam et al. (2019)
Celtidaceae	<i>Celtis integrifolia</i>	Garba et al. (2019)
Cochlospermaceae	<i>Crotalaria rotusa</i>	Garba et al. (2019)
	<i>Pilostigma reticulatum</i>	Garba et al. (2019)
Combretaceae	<i>Anogeissus leiocarpa</i>	Dassou et al. (2012); Ogni et al. (2014); Garba et al. (2019);
	<i>Combretum micranthum</i>	Garba et al. (2019)
	<i>Guiera senegalensis</i>	Garba et al. (2019)
	<i>Pteleopsis suberosa</i>	Dassou et al. (2012); Ogni et al. (2014)
	<i>Terminalia avicennioides</i>	Dassou et al. (2012);
	<i>Terminalia macroptera</i>	Dassou et al. (2012); Ogni et al. (2014)
Convolvulaceae	<i>Ipomoea asarifolia</i>	Dassou et al. (2012); Ogni et al. (2014)
	<i>Ipomoea batatas</i>	Dassou et al. (2012)
	<i>Ipomoea eriocarpa</i>	Dassou et al. (2012)
Crassulaceae	<i>Kalanchoe crenata</i>	Dassou et al. (2012)
	<i>Adenopus breviflorus</i>	Dassou et al. (2012)
Cucurbitaceae	<i>Cucumis melo</i>	Attindehou et al. (2012)
	<i>Momordica charantia</i>	Dassou et al. (2012)
	<i>Croton zambesicus</i>	Garba et al. (2019)
Euphorbiaceae	<i>Euphorbia kamerunica</i>	Dassou et al. (2012); Ogni et al. (2014)
	<i>Jatropha curcas</i>	Attindehou et al. (2012)
	<i>Phyllanthus amarus</i>	Attindehou et al. (2012)
	<i>Acacia albida</i>	Garba et al. (2019)
	<i>Acacia nilotica</i>	Garba et al. (2019)

Fabaceae	<i>Cajanus cajan</i>	Attindehou et al. (2012)
	<i>Gliricidia sepium</i>	Attindehou et al. (2012)
	<i>Tephrosia purpurea</i>	Garba et al. (2019)
Lamiaceae	<i>Ocimum canum</i>	Attindehou et al. (2012)
	<i>Ocimum gratissimum</i>	Attindehou et al. (2012)
Leguminosae	<i>Caesalpinia bonduc</i>	Dassou et al. (2012)
	<i>Caesalpinia pulcherrima</i>	Dassou et al. (2012)
	<i>Detarium microcarpum</i>	Dassou et al. (2012)
	<i>Isobertinia doka</i>	Dassou et al. (2012); Ogni et al. (2014)
	<i>Prosopis africana</i>	Dassou et al. 2012; (Garba et al. 2019)
Lythraceae	<i>Lawsonia inermis</i>	Garba et al. (2019);
Meliaceae	<i>Khaya senegalensis</i>	Dassou et al. (2012); Garba et al. (2019)
	<i>Trichilia prieuriana</i>	Attindehou et al. (2012)
	<i>Azadirachta indica</i>	Dassou et al. (2012); Attindehou et al. (2012); Ogni et al. (2014);
Menispermaceae	<i>Rhigiocarya racemifera</i>	Dassou et al. (2012); Ogni et al. (2014)
Moraceae	<i>Ficus unibellata</i>	Attindehou et al. (2012)
Moringaceae	<i>Moringa oleifera</i>	Dassou et al. (2012); Attindehou et al. (2012); Sidi et al. (2017a); Ogni et al. (2014)
Myrtaceae	<i>Psidium guajava</i>	Dassou et al. (2012); Attindehou et al. (2012); Ogni et al. (2014); Silva et al. (2020)
Olacaceae	<i>Ximenia americana</i>	Dassou et al. (2012)
Phytolaccaceae	<i>Petiveria alliacea</i>	Attindehou et al. (2012)
	<i>Eragrostis tremula</i>	Garba et al. (2019)
Poaceae	<i>Zea mays</i>	Kpabi et al. (2020)
Polygalaceae	<i>Securidaca longepedunculata</i>	Garba et al. (2019)
Rhamnaceae	<i>Ziziphus mauritiana</i>	Garba et al. (2019)
Rubiaceae	<i>Gardenia erubescens</i>	Kabore et al. (2007)
	<i>Macrosphyra longistyla</i>	Dassou et al. (2012)
	<i>Sarcocephalus latifolius</i>	Dassou et al. (2012); Ogni et al. (2014)
	<i>Spermacoce stachydea</i>	Garba et al. (2019)
Rutaceae	<i>Citrus limon</i>	Dassou et al. (2012)
	<i>Citrus sinensis</i>	Ogni et al. (2014)
	<i>Harrisonia abyssinica</i>	Attindehou et al. (2012)
Salvadoraceae	<i>Salvadora persica</i>	Garba et al. (2019)
Scrophulariaceae	<i>Striga hermonthica</i>	Garba et al. (2019)
Zygophyllaceae	<i>Balanites aegyptiaca</i>	Garba et al. (2019)

1.4. Secondary metabolites responsible for the anthelmintic activity of medicinal plants

The presence of bioactive substances such as flavonoids, alkaloids, tannins and essential oils confer anthelmintic properties to medicinal plants (Williams et al. 2014; Azando et al. 2017; Olounladé et al. 2017; Saha and Lachance 2020; Castro et al. 2021), Tannins reportedly inhibit larval migration and egg hatching (Olounladé et al. 2017; Tianhoun et al. 2020; Tchétan et al. 2020). Tannins will bind to free proteins in the cuticle which will then result in physiological alteration of motility, nutrition, absorption and reproduction (Githiori et al. 2006). Fig. 2 describes the mechanisms and sites of action of secondary metabolites of medicinal plants with anthelmintic properties.

1.5. Factors that may influence the anthelmintic activity of medicinal plants

The results obtained on the effectiveness of plant extracts on gastrointestinal parasites, depend on several parameters such as: the species part used; the type of extract; the dose of extract applied, and the parasite used (life stage). According to the work of Akouedegni et al. (2019), the ethanolic extract of *S. monbin* leaves is more effective on larval motility than the methanolic extract. Similar observations are reported by Ademola et al. (2005), according to which the aqueous extract of the same *S. monbin* plant is more effective on larvae than the ethanolic extract. The acetone extract of *P. biglobossa* is more effective on larval migration than methanolic extract (Dedehou et al. 2014); the acetonic extract of *Cassia alata* is more effective than the aqueous extract on egg hatching (Tianhoun et al. 2020). This difference in the activity of one extract of the same plant compared to another would be related to the presence or absence of certain secondary metabolites in one extract compared to the other (Tianhoun et al. 2020). Indeed, depending on affinities, a solvent has the potential to extract in majority certain secondary metabolites. However, the production of secondary metabolites in plant species is correlated to geographical, climatic and genetic factors of the species (Maiga et al. 2020). Essential oils extracted from medicinal plants are also subject to much variation in their chemical compositions. They can vary depending on the geographical site,

time of day, season and genetic factors specific to the species itself. For a better use of anthelmintic medicinal plants, it is therefore necessary to master the factors of variation of secondary metabolites responsible for anthelmintic activities. It is also important to focus on the bioavailability of the species for a better exploitation of these medicinal plants.




Fig. 2: Potential mechanisms of action of secondary metabolites (essential oil; tannins flavonoids, alkaloids, phenol) from medicinal plants on gastrointestinal parasites of small ruminants.

Conclusion: Through previous studies, medicinal plants have demonstrated their ability to be used in the control of gastrointestinal parasitosis in small ruminants. However, the anthelmintic activities of these plants depend on several factors that must be controlled. The secondary metabolites: flavonoids, alkaloids, tannins and essential oils responsible for the anthelmintic activity of medicinal plants are subject to variations according to geographical and intrinsic factors of the plants. Also, the secondary metabolites present in a given medicinal plant extract depend on the solvent used for extraction. Anthelmintic medicinal plants (or their extracts) do not have the same efficacy on parasitic nematodes, and small ruminants and even humans are often confronted with polyparasitism. In short, anthelmintic medicinal plants are good candidates in the fight against intestinal parasitosis.

ORCID

Lisette Degla

<https://orcid.org/0000-0002-3304-3459>

Julienne Kuiseu

<https://orcid.org/0000-0002-4574-9184>

Pascal Abiodoun Olounlade

<https://orcid.org/0000-0003-1173-3536>

Sabbas Attindehou

<https://orcid.org/0000-0001-8172-0138>

Mawulé Sylvie Hounzangbe-Adote

<https://orcid.org/0000-0001-6845-3477>

Patrick A. Eдорh

<https://orcid.org/0000-0001-8761-2461>

Latifou Lagnika

<https://orcid.org/0000-0003-2942-9372>

REFERENCES

- Ademola IO, Fagbemi BO and Idowu SO, 2005. Anthelmintic activity of extracts of *Spondias mombin* against gastrointestinal nematode of sheep: Studies *in vitro* and *in vivo*. *Tropical Animal Health and Production* 37: 223-235. <https://doi.org/10.1023/B:TROP.0000049296.47350.80>
- Aguerre S, 2019. Résistance génétique aux nématodes gastro-intestinaux chez les ovins: évaluation des stratégies de sélection et de leur impact à l'échelle de l'élevage. Thèse de Doctorat de l'Université de Toulouse, pp: 172.
- Ahmed AH, Ejo M, Feyera T, Regassa D, Mammed B and Huluka SA, 2020. *In Vitro* Anthelmintic Activity of Crude Extracts of *Artemisia herba-alba* and *Punica granatum* against *Haemonchus contortus*. *Journal of Parasitology Research* 2020: Article # 4950196. <https://doi.org/10.1155/2020/4950196>
- Akoto CO, Acheampong A, Boakye YD, Akwata D and Okine M, 2019. *In vitro* anthelmintic, antimicrobial and antioxidant activities and FTIR analysis of extracts of *Alchornea cordifolia* leaves. *Journal of Pharmacognosy and Phytochemistry* 8: 2432-2442.
- Akouedegni CG, Daga FD, Olounlade PA, Allowanou GO, Ahoussi E, Tamboura HH and Hounzangbe-Adote MS, 2019. Evaluation *In vitro* et *in vivo* des propriétés anthelminthiques de feuilles de *Spondias mombin* sur *Haemonchus contortus* des ovins Djallonke. *Agronomie Africaine* 31: 213-222.

- Alowanou GG, Azando EVB, Adenilé AD, Koudandé DO, Chrysostome CAM and Hounzangbé-Adoté SM, 2019a. Evaluation of the *in vivo* anthelmintic properties of *Mitragyna inermis* (Willd.) as a livestock dewormer against parasitic hematophagous worm *Haemonchus contortus* infections in different breeds of lambs. *Tropical Animal Health and Production* 52: 9-19. <https://doi.org/10.1007/s11250-019-02014-w>
- Alowanou GG, Olounladé PA, Akouèdegne GC, Faihun AML, Koudandé DO and Hounzangbé-Adoté S, 2019b. *In vitro* anthelmintic effects of *Bridelia ferruginea*, *Combretum glutinosum*, and *Mitragyna inermis* leaf extracts on *Haemonchus contortus*, an abomasal nematode of small ruminants. *Parasitology Research* 118: 15-23. <https://doi.org/10.1007/s00436-019-06262-5>
- Alowanou GG, Olounladé AP, Koudandé OD, Babatoundé S and Hounzangbé-Adoté MS, 2015. Effets de la digestion dans le rumen sur les propriétés anthelminthiques de *Bridelia ferruginea* (Benth.), *Mitragyna inermis* (Willd.) Kuntze et *Combretum glutinosum* (Perr. ex DC.). *Science de la Vie, de la Terre et Agronomie* 3: 50-56.
- Attindehou S, Salifou S, Biaoou CF, Gbati OB, Diaye AN and Pangui LJ, 2012. Epidemiology of haemonchosis in sheep and goats in Benin. *Journal of Parasitology and Vector Biology* 42: 20-24. <https://doi.org/10.58897/JPV12.012>
- Azando EVB, Olounlade AP, Hounzangbe-Adote MS and Hoste H, 2011. Effets anthelminthiques *in vivo* de la poudre de feuilles de *Zanthoxylum zanthoxyloides* et de *Newbouldia laevis* sur les nématodes parasites gastro-intestinaux des chevreaux Djallonké. *International Journal of Biologie and Chemical Sciences* 5: 1054-1062.
- Azando EVB, Olounlade AP, Hounzangbe-Adote MS, Tam Ha TB, Fabre N and Valentin A, 2017. Contrôle des parasitoses gastro-intestinales ovines par l'huile essentielle de *Zanthoxylum zanthoxyloides* (*Fagara zanthoxyloides*). *Revue de Médecine Vétérinaire* 168: 205-212.
- Castro LM, Pinto NB, Moura MQ, Villela MM, Capella GA, Freitag RA and Berne MEA, 2021. Anthelmintic action of the *Anethum graveolens* essential oil on *Haemonchus contortus* eggs and larvae. *Brazilian Journal of Biology* 81: 183-188. <https://doi.org/10.1590/1519-6984.225856>
- Coyne MJ and Smith G, 1992. The development and mortality of the free-living stages of *Haemonchus contortus* in laboratory culture. *International Journal of Parasitology* 22: 641-650.
- Dassou GH, Adomou AC, Yédomonhan H, Ogni AC, Tossou GM, Dougnon JT and Akoègninou A, 2015. Flore médicinale utilisée dans le traitement des maladies et symptômes animaux au Bénin. *Journal of Animal and Plant Sciences* 26: 4036-4057.
- Dassou HG, Ogni CA, Yédomonhan H, Adomou AC Tossou M, Dougnon JT and Akoègninou A, 2015. Facteurs socioculturels et environnementaux déterminant la connaissance ethnovétérinaire au Bénin. *Afrique Science* 11: 335-360.
- Dedehou VFGN, Olounlade PA, Adenilé AD, Azando EVB, Alowanou GG, Daga FD and Hounzangbe-Aoté MS, 2014. Effets *In vitro* des feuilles de *Pterocarpus erinaceus* et des cosses de fruits de *Parkia biglobosa* sur deux stades du cycle et développement de *Haemonchus contortus* nématode parasite gastro intestinal de petits ruminants. *Journal for Animal and Plant Sciences* 22: 3368-3378.
- Emery DL, Hunt PW and Le Jambre LF, 2016. *Haemonchus contortus* : the then and now, and where to from here? *International Journal for Parasitology* 46: 755-769. <https://doi.org/10.1016/j.ijpara.2016.07.001>
- Ferreira LE, Benincasa BI, Fachin A, Franca SC, Contini SSHT, Chagas ACS and Belebani RO, 2016. *Thymus vulgaris* L. essential oil and its main component thymol: Anthelmintic effects against *Haemonchus contortus* from sheep. *Veterinary Parasitology* 228: 70-76. <https://doi.org/10.1016/j.vetpar.2016.08.011>
- Furtado LFV, Medeiros CS, Zuccherato LW, Alves WP, de Oliveira VNGM and da Silva VJ, 2019. First identification of the benzimidazole resistance-associated F200Y SNP in the beta-tubulin gene in *Ascaris lumbricoides*. *PLoS One* 14: e0224108. <https://doi.org/10.1371/journal.pone.0224108>
- Garba ARI, Adakal H, AbasseT, Koudouvo K, Karim S, Akourki A, Gbeassor M and Mahamane S, 2019. Etudes ethnobotaniques des plantes utilisées dans le traitement des parasitoses digestives des petits ruminants (ovins) dans le Sud-Ouest du Niger. *International Journal of Biologie and Chemical Sciences* 13: 1534-1546. <https://doi.org/10.4314/ijbcs.v13i3.26>
- Githiori JB, Athanasiadou S and Thamsborg SM, 2006. Use of plants in novel approaches for control of gastrointestinal helminths in livestock with emphasis on small ruminants. *Veterinary Parasitology* 139: 308-320. <https://doi.org/10.1016/j.vetpar.2006.04.021>
- Houngnimassoun HMA, Attindehou S, Salifou S, Koumodji KD and Salifou S, 2017. Effets stronglycidés *In vitro* de l'extrait aqueux de feuilles de *Ficus exasperata* Valh. 1805 (Moraceae). *International Journal of Biological and Chemical Sciences* 11: 1012-1020. <https://doi.org/10.4314/ijbcs.v11i3.6>
- Islam R, Zahra SFT, Sumon SMI, Parvin S, Hasan K, Ahmed M, Siddique AT and Haque T, 2019. Evaluation of anthelmintic activity of ethanolic extracts of *Carica papaya* leaves using *Paramphistomum cervi* and *Haemonchus contortus*. *African Journal of Pharmacy and Pharmacology* 13: 146-150. <https://doi.org/10.5897/AJPP2019.5017>
- Kabore A, Tamboura HH, Belem AMG and Traore A, 2007. Traitements ethno-vétérinaires des parasitoses digestives des petits ruminants dans le plateau central du Burkina Faso. *International Journal of Biological and Chemical Science* 1: 297-304.
- Kpabi I, Agban A, Hoekou Y, Pissang P, Tchacondo T and Batawila K, 2020. Etude ethnobotanique des plantes à activités antiparasitaires utilisées en médecine traditionnelle dans la préfecture de Doufelgou au nord du Togo. *Journal of Applied Biosciences* 148: 15176-15189. <https://doi.org/10.35759/JABs.148.2>
- Kuiseu J, Zinsou FTA, Olounladé PA, Alowanou GG, Adenilé AD, Dansou CC, Hounzangbé-Adoté MS, Babayemi OJ and Edoh PA, 2021. Prevalence, effects and alternative control methods of *Haemonchus contortus* in small ruminants: Review. *Journal of Veterinary Medicine and Animal Health* 13: 84-97. <https://doi.org/10.5897/JVMAH2020.0868>

- Maiga A, Houngrimmassoun HMA, Attindehou S, Houinato M and Salifou S, 2020. Effet vermicide *in vitro* de l'extrait aqueux des feuilles de *Chenopodium ambrosioides* L. 1753 sur *Haemonchus contortus* et *Oesophagostomum colombianum* parasites gastro-intestinaux des petits ruminants. *Journal of Animal and Plant Sciences* 43: 7501-7512. <https://doi.org/10.35759/JAnmPISci.v43-3.6>
- Mesmin KY, Kamelé KY, Gervaises KA, Jules KNG and Adama B, 2018. Activité anthelminthique *in vitro* et teneurs en tanins et flavonoïdes de huit plantes fourragères utilisées en élevage des petits ruminants en Côte d'Ivoire. *European Scientific Journal* 14: 434-449. <https://doi.org/10.19044/esj.2018.v14n15p434>
- Nath S, Pal S, Sanyal P, Roy S, Mandal S and Nayak Y, 2019. Anthelmintic activity of *Curcuma Longa* ethanolic extract against benzimidazole resistant gastrointestinal nematodes in goats. *International Journal of Livestock Research* 9: 117-122. <https://doi.org/10.5455/ijlr.20190209020421>
- Nguessan CB, Koutouan FP, Wandan EN, Yapi YM, N'din NS and Iritié BM, 2017. Évaluation *in vivo* des propriétés anthelminthiques des extraits aqueux d'*Annona senegalensis*. Pers. (Annonaceae) chez les ovins Djallonkés. *Journal of Animal and Plant Sciences* 32: 5111-5119.
- Ogni CA, Kpodekon MT, Dassou HG, Boko CK, Koutinhoun BG, Dougnon JT, Youssao AKI, Yedomonhan H and Akoegninou A, 2014. Inventaire ethno-pharmacologique des plantes utilisées dans le traitement des pathologies parasitaires dans les élevages extensifs et semi-intensifs du Bénin. *International Journal of Biologie and Chemical Sciences* 8: 1089-1102.
- Okombe EV and Pongombo SEW, 2013. Suspicion de la résistance aux benzimidazoles chez les strongles gastro-intestinaux du caprin à Lubumbashi, R.D. Congo. *International Journal of Biologie and Chemical Sciences* 7: 2426-2433.
- Olounladé PA, Azando EVB, Attakpa EY, Gbneou JD, Alowanou GG, Tchétan E, Dansou CC, Hounzangbé-Adoté MS, Gbaguidi F, Moudachirou M, Hoste H and Valentin A, 2017. *In vitro* study on the role of the tannins of *Newbouldia laevis* and *Zanthoxylum zanthoxyloides* on infective larvae of *Trichostrongylus colubriformis*. *African Journal of Agricultural Research* 12: 3513-3519. <https://doi.org/10.5897/AJAR11.1353>
- Olukotun AB, Bello IA and Oyewale OA, 2018. Phytochemical and anthelmintic activity of *Terminalia catappa* (Linn) leaves. *Journal of Applied Sciences and Environmental Management* 22: 1343-1347. <https://doi.org/10.4314/jasem.v22i8.33>
- Rehman T, Iqbal KJ, Anwer A, Abbas RZ, Babar W, Ali A, Khan MK, Javid A, Khan N, Ali HM, Shahzad F, Zaman MA and Sikandar A, 2021. *In vitro* anthelmintic efficacy of *Citrullus colocynthis* (L.) Schrad on *Haemonchus contortus*. *Veterinarski Arhiv* 91: 309-318. <https://doi.org/10.24099/vet.arhiv.1024>
- Saha S and Lachance S, 2020. Effect of essential oils on cattle gastrointestinal nematodes assessed by egg hatch, larval migration and mortality testing. *Journal of Helminthology* 94: e111. <https://doi.org/10.1017/S0022149X19001081>
- Sidi IYMS, Alowanou GG, Tchétan E, Aminou MY, Hounzangbe-Adote MS and Babatoundé S, 2017b. Effets De La Digestion gastrique sur les Propriétés anthelminthiques de *Zanthoxylum Zanthoxyloides*, (Lam.) Zepernick & Timlerto Et De *Newbouldia, Laevis* (P.Beauv.) Sur *Haemonchus Contortus*. *European Scientific Journal* 13: 204-216. <http://dx.doi.org/10.19044/esj.2017.v13n24p204>
- Sidi IYMS, Olounladé PA, Yaoitcha A, Dedehou VFGN, AloXwanou GG, Azando EVB and Hounzangbe-Adote MS, 2017b. Principales espèces médicinales utilisées en médecine vétérinaire au Bénin: disponibilité et caractéristiques dendrométriques. *Bulletin of Animal Health and Production in Africa* 65: 209-220.
- Silva LP, Debiage RR, Bronzel-Júnior JL, Da Silva RMG and Mello-Peixoto ECT, 2020. *In vitro* anthelmintic activity of *Psidium guajava* hydroalcoholic extract against gastrointestinal sheep nematodes. *Anais Academia Brasileria Ciencias* 92: 5-15.
- Sobhy H, AboElnaga TR, Behour TS and Razin EA, 2021. *In vitro* trypanocidal activity of essential oils of some plants against *Trypanosoma evansi*. *International Journal of Veterinary Science* 10: 191-195. <https://doi.org/10.47278/journal.ijvs/2021.043>
- Tchetan E, Azando EVB, Olounladé PA, Alowanou GG and Hounzangbé-Adoté SM, 2020. *In vitro* effects of tannin and extracts of *Bridelia ferruginea* and *Mitragyna inermis* on the exsheathment of infective larvae of *Haemonchus contortus*. *International Journal of Veterinary Science and Medicine* 8: 93-99. <https://doi.org/10.1080/23144599.2020.1837056>
- Tianhoun DF, Meda N-TR, Konate A, Kabore A, Tamboura HH and Belem AMG, 2020. Phytochemical screening and *in vitro* anthelmintic activity of *Cassia alata* (L) Roxb. on *Haemonchus contortus* of small ruminants in Burkina Faso. *Journal of Advances in Parasitology* 7: 14-19. <http://doi.org/10.17582/journal.jap/2020/7.3.14.19>
- Toklo PM, Ladekan Yai EC, Sakirigui A, Alowanou GG, Assogba MF, Challaton KP, Hounzangbe-Adote S and Gbenou JD, 2021. Antioxidant and antiparasitic activities on *Trichostrongylus colubriformis* of the aqueous extract of fresh leaves of *Mitragyna inermis* (Willd.) Kuntze (Rubiaceae). *Pharmaceutical and Chemical Journal* 8: 120-128.
- Williams AR, Frygas C, Ramsay A, Mueller-Harvey I and Thamsborg SM, 2014. Direct anthelmintic effects of condensed tannins from diverse plant sources against *Ascaris suum*. *PLoS One* 9: e97053. <https://doi.org/10.1371/journal.pone.0097053>
- Yongwa G, BFNF Ngnodandi, Ndjonka D and Saotoing P, 2020. *In vitro* Anthelmintic Activity of aqueous and ethanolic extract of *Senna italica* (Caesalpinaceae) on three-stages of *Haemonchus contortus*. *Journal of Pharmaceutical Research International* 32: 25-34. <https://doi.org/10.9734/jpri/2020/v32i330411>
- Zangueu CB, Olounladé AP, Ossokomack M, Djouatsa YNN, Alowanou GG, Azebaze AGB, Llorent-Martínez EJ, de Córdova MLF, Dongmo AB and Hounzangbe-Adote MS, 2018. *In vitro* effects of aqueous extract from *Maytenus senegalensis* (Lam.) Exell stem bark on egg hatching, larval migration and adult worms of *Haemonchus contortus*. *BMC Veterinary Research* 14: 147. <https://doi.org/10.1186/s12917-018-1475-3>
- Zinsou AL, 2015. Evaluation *ex vivo* de l'extrait aqueux de *Pleiocarpa pycnantha* (k.schum) sur *Haemonchus contortus*, nématode parasite gastro-intestinale des petits ruminants. Mémoire pour l'obtention du diplôme de master en physiologie et applications soutenu à l'Université d'Abomey-Calavi, pp: 63.