

## BIO-EFFICACY OF GROUNDED *SESAMUM RADIATUM* AND *PSORALEA CORYLITOLIA* ON EMERGENCE OF *ANOPHELES GAMBIAE* AND *CULEX QUINQUEFASCIATUS*

Ojianwuna CC<sup>ORCID</sup>, Enwemiwe VN<sup>ORCID</sup>\*, Ojo MG<sup>ORCID</sup> and Oborayiruvbe ET<sup>ORCID</sup>

Department of Animal and Environmental Biology, Faculty of Sciences,  
Delta State University Abraka, Delta State, Nigeria

\*Corresponding author: [enwemiwevector@gmail.com](mailto:enwemiwevector@gmail.com)

### ABSTRACT

Mosquito larvae are highly distributed species in tropical regions where breeding sites are diverse and causing a number of public health diseases, when they emerge. This study tested the efficacy of two grounded insecticidal plant powders against the emergence of *Anopheles* and *Culex* species. Larvae of mosquitoes were collected using 350mL deep ladles and scooping spoon from Ethiope East LGA, Delta State, Nigeria. The mosquito larvae were exposed to 0.5, 1.0, and 1.5g of *Psoralea corylitolia* and *Sesamum radiatum* in 100mL of water. This corresponded to 0.005, 0.01, and 0.015g/mL concentration of experimental set up respectively. Mosquitoes were exposed to these treatments in 20 larvae per concentration in quadruplicates. Mean mortality and time of mortality were recorded. Probit model was used to determine  $LT_{50}$  and  $LT_{95}$ . Mortality of larvae increased as concentration and time increased. Highest mortality was recorded in *Culex* mosquitoes exposed to 1.5g of *Psoralea corylitolia* and was closely followed by *Anopheles* mosquitoes exposed to same concentration of *Sesamum radiatum*. The differences between the mortalities were highly significant ( $P < 0.00001$ ). Emergence depreciated as the concentration increased with high mortality especially with 0.5g of *Psoralea corylitolia* exposed to *Anopheles* mosquito larvae. The insecticidal test plants did not determine the sex of mosquitoes and it was not significant ( $F_{sex\ ratio} = 1.127$ ;  $P = 0.357$ ). Physicochemical parameters supported the mortality of both larvae exposed to *Sesamum radiatum* from 30 minutes till end of the post experimental exposure. Lethal time for 50% of mosquitoes ranged from 31 to 5858min whereas  $LT_{95}$  ranged from 1435 to 10218min for the two insecticidal test plants respectively. There is need for more study with the trial of oil of *Sesamum radiatum* and the determination of the gas chromatograph of this novel plant to ascertain the active components.

**Keywords:** Bio-efficacy, emergence, mosquitoes, *Sesamum radiatum*, *Psoralea corylitolia*.

---

Article History (2022-12139) || Received: 25 Dec 2022 || Revised: 27 Dec 2022 || Accepted: 28 Dec 2022 || Published Online: 31 Dec 2022

---

### 1. INTRODUCTION

Mosquitoes are highly abundant vectors that cause several diseases of increasing public health concern in humans and animals globally (WHO 2017). Unlike vectors of other infectious diseases, mosquitoes have been spotted as the most important due to the role they play in disease transmissions (Becker et al. 2020). Some of these diseases including dengue fever, malaria, filariasis, arboviruses, west Nile virus, Japanese encephalitis, yellow fever, chikungunya and so many of them (Chancey et al. 2015; Dad et al. 2019). Amongst these diseases, malaria stands out as the major human disease in Africa and Nigeria, with about 97% of Nigerian population at risk (Anonymous 2014; Nwaneli et al. 2020). Apart from the fact that mosquitoes transmit several diseases, their bites can in rare cases cause associated syndrome with swellings, sore, reddish spots that are itchy and presented with pain (Lopez 2022). Approximately 3500 species of mosquitoes are known globally with the exception of the polar regions (CDC 2020). Mosquitoes are spread in various parts of the world. There are about 100 mosquito species native to Europe, over 870 species to South-east Asia, and Africa is considered home for diversities of species due to the warm climate that favors life stage development (Lai et al. 2018; Kampen and Walther 2018; Trájer and Padisák 2019; Steinbrink et al. 2022). The prominent species causing mosquito-borne diseases in Nigeria are common in the genera *Anopheles*, *Aedes* and *Culex*. Mosquito-biting activities and diseases cause economic losses amounting to over one billion dollars in vector and disease management (Hamlet et al. 2022).

In order to reduce the burdens of continuous budgeting for disease management, it is important to control mosquitoes through the use of insecticides in the form of larvicides or adulticides, manipulation of physicochemical

characteristics of breeding habitat, physical controls in the use of traps, biological control in the use of *Bacillus thuringiensis* (BTI), larvivorous fishes and plant materials. The use of insecticides, either chemical or plant-extracted, is recommended in the forms of treated bed nets, treated clothing, personal protection by rubbing repellents and spraying indoor of houses (Kimani et al. 2006; Farag et al. 2021). Insecticide treated nets and materials confer 75% protection against mosquito bites and guaranteed the reduction of deaths in vulnerable children under the age of 5 by 20% (CDC 2019). The spraying of insecticide indoors accounted for 60% of insecticide usage in vector control. The issues of resistance in adult vectors have diverted attention towards the use of growth regulators, bacterial and plant extracts in the control of immature stages of mosquitoes in their breeding habitats (Mahran 2022). Oil film made by the oil extracts of *Azadirachta indica* and many other plants was successful in killing immature stages of *Anopheles*, *Culex*, and *Aedes* mosquitoes (Ayinde et al. 2020). Plant-based oil has been proven to be environmentally friendly, effective and poses lesser threat to human health and the environment. This is for their active components that are target specific on the vectors. There is no information on the biological efficacy of these plants on the emergence of *Anopheles* and *Culex* mosquitoes. Therefore, this study was conducted.

## 2. MATERIALS AND METHODS

### 2.1. Mosquito Collection

Immature stages of mosquito for this study were collected using 350mL dip ladles and scooping spoons from potential breeding habitat including ponds, tire marks, and puddles in Ethiopia East Local Government Area, Delta State. These immature stages were transferred to the Entomology Laboratory in transparent buckets. The larvae of *Anopheles* were distinguished from *Culex* using the resting position on water surfaces. *Anopheles* larvae lie 180° to the water surface and *Culex*, at 45°.

### 2.2. Plant Source and Preparation

The plants, *Psoralea corilifolia* and *Sesamum radiatum*, for this experiment were sourced from the environment. These plants were air dried at room temperature and grounded into powder and stored in airtight containers which were well labelled.

### 2.3. Experimental Design

Complete randomized exposure was used in this study. Twenty (20) larval stage three (L<sub>3</sub>) of *Anopheles* and *Culex* were sorted into 1000mL capacity glass exposure vials and assigned to four replicates to various concentrations of 0.5, 1.0, and 1.5g, respectively in 100mL of water. Readings for acute and chronic toxicity were taken. Acute toxicity was taken at 10, 20, 30, 40, 50, and 60min. Chronic toxicity was taken from 6-72hrs. Plastic pipette was used to prickle the larvae in the case where larvae show sign of death. Larvae in various experimental set up were fed biscuit and yeast powder.

### 2.4. Physicochemical Analysis

Multiplex water analysis and pH device was used to take readings for pH, water conductivity, salinity, and temperature of water in the set up.

### 2.5. Data Analysis

Data were entered into Microsoft Excel Spreadsheet and checked thoroughly. Descriptive statistics were used in the result presentation. Analysis of Variance (ANOVA) test was used to test significance at P=0.05. Mortality was correlated with physicochemical parameters of water to check for correspondence.

## 3. RESULTS

### 3.1. Mortality of Mosquitoes

Mean mortality of mosquito larvae exposed to two insecticidal test plants in Ethiopia East LGA, Delta State, Nigeria is shown in Table 1. The mortality of larvae increased as concentration increased with the two mosquitoes, but a slight decline was observed in *Culex* exposed to 1.0g of *Psoralea corylitolia*. The highest mortality was recorded in *Culex* mosquitoes exposed to 1.5g of *Psoralea corylitolia*. This was closely followed by *Anopheles* mosquitoes exposed to same concentration of *Sesamum radiatum*. The differences between the mortalities of mosquito larvae were highly significant (P=0.0000).


### 3.2. Mortality

The percentage time mortality of *Anopheles* and *Culex* mosquitoes to two insecticidal test plants is shown in Fig. 1 and 2. Generally, mortality increased with time as concentration increased. 1.5g of *Psoralea corylitolia*


caused higher mortality in *Culex* mosquito larvae than in other concentrations and mosquito. *Psoralea corylitolia* did not cause acute toxicity in less than 1 hour of exposure. The optimum concentration (1.5g) of *Sesamum radiatum* caused high chronic mortality in both *Anopheles* and *Culex*. Other concentrations did too. There was significant mortalities with the use of insecticidal plants against the emergence of *Anopheles* and *Culex*.

**Table I:** Mean mortality of mosquito larvae exposed to two insecticidal test plants in Ethiope East LGA, Delta State, Nigeria

Plants	Mosquito	Conc. (gram)	Mean±SE	F-ANOVA	P-value
<i>Psoralea corylitolia</i>	<i>Anopheles</i>	0.5	5.64±3.65	35.98	0.00000
	<i>Culex</i>	0.5	22.55±4.02		
	<i>Anopheles</i>	1.0	7.64±4.59		
	<i>Culex</i>	1.0	20.73±4.43		
	<i>Anopheles</i>	1.5	10.36±5.86		
	<i>Culex</i>	1.5	47.46±7.78		
<i>Sesamum radiatum</i>	<i>Anopheles</i>	0.5	22.91±7.82		
	<i>Culex</i>	0.5	26.91±9.62		
	<i>Anopheles</i>	1.0	31.46±8.50		
	<i>Culex</i>	1.0	33.27±8.50		
	<i>Anopheles</i>	1.5	45.64±9.18		
	<i>Culex</i>	1.5	43.46±7.63		




**Fig. 1:** Percentage time mortality of *Anopheles* and *Culex* mosquitoes to *Psoralea corylitolia* (F=19.57; P=0.000001).



**Fig. 2:** Percentage time mortality of *Anopheles* and *Culex* mosquitoes to *Sesamum radiatum* (F=56.35; P=0.00001).

Some mosquitoes emerged in various concentrations. The percentage emergence of *Anopheles* and *Culex* mosquitoes to two insecticidal test plants is shown in Fig. 3 and 4. Emergence depreciated as the concentration increased with high mortality. The highest emergence in the *Psoralea corylitolia* was recorded with 0.5g of *Psoralea corylitolia* exposed to *Anopheles* mosquito larvae. Emergence in the *Sesamum radiatum* was higher in 0.5 and 1.0g of the plant exposed to *Anopheles* and *Culex* mosquitoes. There was non-significance in the emergence of *Anopheles* and *Culex* mosquitoes to the insecticidal plants (P>0.05).



**Fig. 3.** Percentage emergence of *Anopheles* and *Culex* mosquitoes to *Psoralea corylitolia* (F=2.107; P=0.135).

**Fig. 4:** Percentage emergence of *Anopheles* and *Culex* mosquitoes to *Sesamum radiatum* (F=0.161; P=0.972).

### 3.3. Sex of Emerged Mosquitoes

Sex ratio of mosquito emergence after exposure to two insecticidal test plants with time is shown in Table 2. The insecticidal test plants did not determine the sex of mosquitoes. There were more emerged female mosquitoes (n=146) than males (n=98). Male and female *Anopheles* were 66 and 84, respectively after exposure to plants. Similarly, male and female *Culex* were 32 and 62, respectively after exposure to plants. The difference in sex of mosquitoes exposed to the plants was not significant ( $F_{sex\ ratio}=1.127$ ;  $P=0.357$ ).

### 3.4. Correspondence Analysis


Correspondence analysis between physicochemical parameters and mortality of mosquito larvae to two insecticidal test plants in Ethiopie East LGA, Delta State, Nigeria is shown Fig. 5. It appeared that all parameters except salt supported the mortality of *Culex* mosquito larvae more than in *Anopheles* mosquito larvae when exposed to powder of *Psoralea corylitolia*. Similarly, all parameters supported the mortality of both larvae exposed to *Sesamum radiatum* from 30 minutes to the end of the post experimental exposure.

### 3.5. Lethal Time Toxicity

Lethal time model of mosquito larvae exposed to two insecticidal test plants in Ethiopie East LGA, Delta State, Nigeria is shown in Table 3. Lethal time for 50% of mosquitoes ranged from 31 to 5858min whereas  $LT_{95}$  ranged from 1435 to 10218min for the two insecticidal test plants respectively. *Culex* mosquitoes exposed to 1.5g of *Psoralea corylitolia* recorded the lowest lethal time and this followed by *Anopheles* mosquitoes exposed to same concentration of *Sesamum radiatum*. The lethal time model of the mosquitoes followed a well fitted curve as observed in the Pearson goodness of fit and  $r^2$  (Table 3).

**Table 2:** Sex ratio of mosquito emergence after exposure to two insecticidal test plants with time.

Plant	Mosquito	Conc. (g)	Sex	24hrs (%)	48hrs (%)	72hrs (%)
Psoralea corylitolia	Anopheles	0.5	♂	14 (17.5)	6 (7.5)	6 (7.5)
			♀	12 (15.0)	8 (10.0)	2 (2.5)
		1.0	♂	10 (12.5)	4 (5.0)	2 (2.5)
			♀	10 (12.5)	8 (10.0)	6 (7.5)
		0.5	♂	14 (17.5)	0 (0.0)	0 (0.0)
			♀	8 (10.0)	6 (7.5)	0 (0.0)
	Culex	0.5	♂	12 (15.0)	0 (0.0)	2 (2.5)
			♀	8 (10.0)	8 (10.0)	6 (7.5)
		1.0	♂	2 (2.5)	6 (7.5)	0 (0.0)
			♀	12 (15.0)	8 (10.0)	0 (0.0)
		0.5	♂	2 (2.5)	0 (0.0)	0 (0.0)
			♀	0 (0.0)	0 (0.0)	0 (0.0)
Sesamum radiatum	Anopheles	0.5	♂	0 (0.0)	0 (0.0)	2 (2.5)
			♀	4 (5.0)	0 (0.0)	6 (7.5)
		1.0	♂	2 (2.5)	4 (5.0)	0 (0.0)
			♀	2 (2.5)	4 (5.0)	0 (0.0)
		0.5	♂	4 (5.0)	0 (0.0)	0 (0.0)
			♀	2 (2.5)	6 (7.5)	0 (0.0)
	Culex	0.5	♂	4 (5.0)	0 (0.0)	0 (0.0)
			♀	4 (5.0)	0 (0.0)	0 (0.0)
		1.0	♂	2 (2.5)	0 (0.0)	0 (0.0)
			♀	6 (7.5)	4 (5.0)	2 (2.5)
		0.5	♂	2 (2.5)	0 (0.0)	0 (0.0)
			♀	4 (5.0)	0 (0.0)	0 (0.0)



**Fig. 5:** Correspondence analysis between physiochemical parameters and mortality of mosquito larvae to two insecticidal test plants in Ethiope East LGA, Delta State, Nigeria.

#### 4. DISCUSSION

This study determined the toxicity of two insecticidal test plants against the emergence of *Anopheles* and *Culex* in Ethiope East LGA, Delta State, Nigeria. This was done to ascertain their biological potency due to the urgent need for toxic materials with ecological and environmentally friendly potential. Different plant substances have been tested for their insecticidal potentials on various insects (Ojjanwuna et al. 2011; Ojjanwuna and Umoru 2011; Ojjanwuna and Enwemiwe 2020b). Even some pyrethroids have been tried on adult mosquitoes (Ojjanwuna and Enwemiwe 2020a). Most larvicidal agents of plant origins are actively toxic against the three major mosquitoes causing disease transmission in Nigeria (Ojjanwuna et al. 2021). Amongst these are the effects of leaf extracts from

**Table 3:** Lethal time model of mosquito larvae exposed to two insecticidal test plants in Ethiopia East LGA, Delta State, Nigeria

Plant Powders	Mosquito Larvae	Conc. (g)	Regression Line	Pearson $\chi^2$ Goodness of Fit ( $r^2$ )	LT <sub>50</sub> (mins)	LT <sub>95</sub> (mins)
<i>Psoralea corylitolia</i>	<i>Anopheles</i>	0.5	$y = 0.010x - 1.98$	9.362 (0.907)	5858	10218
		1.0	$y = 0.013x - 2.01$	11.56 (0.937)	4318	7200
		1.5	$y = 0.017x - 1.99$	14.762 (0.960)	2715	6341
	<i>Culex</i>	0.5	$y = 0.010x - 19.01$	5.496 (0.770)	1536	5759
		1.0	$y = 0.011x + 16.17$	4.763 (0.716)	2859	4312
		1.5	$y = 0.017x + 44.53$	3.222 (0.536)	31	1435
<i>Sesamum radiatum</i>	<i>Anopheles</i>	0.5	$y = 0.021x + 9.50$	8.387 (0.887)	1348	4398
		1.0	$y = 0.020x + 21.46$	4.110 (0.653)	419	4358
		1.5	$y = 0.016x + 42.46$	2.317 (0.374)	56	4327
	<i>Culex</i>	0.5	$y = 0.023x + 12.63$	4.640 (0.705)	295	4331
		1.0	$y = 0.019x + 24.30$	3.773 (0.613)	196	4328
		1.5	$y = 0.016x + 39.65$	3.010 (0.502)	199	4329

**Note:** Lethal time of 50% and 95% (LT<sub>50</sub> and LT<sub>95</sub>) are in minutes; P>0.05 suggests a well-fitting model, P<0.05 suggests an invalid model population.

three asteraceous plants which caused complete or partial mortality. Petroleum products have equally been tried against these mosquitoes (Ekedo et al. 2019; Ojjanwuna et al. 2021b). Plants have been applied as topical treatment for other insect causing disease lesions either as extracted oils or grounded with positive results (Enwemiwe et al. 2020, Enwemiwe et al. 2021). Alam et al. (2018) revealed the toxic potential of *Psoralea corylifolia* in insect studies. Apart from this, plants have several potentials in life applications. Even the toxicity of *Sesamum indicum*, a relative of the plant under study has been tried on *Culex* species.

In this present study, the mortality of larvae increased as concentration increased with the two mosquitoes, but a slight decline was observed in *Culex* exposed to 1.0g of *Psoralea corylitolia*. Mortality was highly recorded in *Culex* mosquitoes exposed to 1.5g of *Psoralea corylitolia* and this was closely followed by *Anopheles* mosquitoes exposed to same concentration of *Sesamum radiatum*. The result of this study corresponded with the findings of Ojjanwuna et al. (2021) where complete mortality was recorded with optimum concentration of *Bidens pilosa*, and *Spilanthes filicaulis*. The mortality was not in agreement with those reported in other insects (Ojjanwuna et al. 2011; Ojjanwuna and Enwemiwe 2020b; Ojjanwuna and Enwemiwe 2021).

Mortality equally increased with time as concentrations of test plants increased. *Psoralea corylitolia* in 1.5g caused higher mortality in *Culex* mosquito larvae than in other concentrations and mosquito. They did not cause acute toxicity in less than 1 hour of exposure. The optimum concentration (1.5g) of *Sesamum radiatum* caused higher chronic mortality in both *Anopheles* and *Culex*. In the findings of the present study, plant powders did not cause acute toxicity in reduced time. It took longer time than possible as compared to the study of Ojjanwuna et al. (2021). Emergence depreciated as the concentration increased with high mortality. The highest emergence in the *Psoralea corylitolia* was recorded with 0.5g of *Psoralea corylitolia* exposed to *Anopheles* mosquito larvae. Emergence in the *Sesamum radiatum* was higher in 0.5 and 1.0g of the plant exposed to *Anopheles* and *Culex* mosquitoes. Emergence inhibition was not determined in this study. To considerable extent, the plant powders did not suppress adult mosquito emergence as reported in the study of Ojjanwuna and Enwemiwe (2022). Insecticidal test plants under study did not determine the sex of mosquitoes. There were more emerged female mosquitoes than males. Most studies do not consider the sex of mosquitoes after emergence unlike in the study of Ojjanwuna and Enwemiwe (2020) where adult mosquitoes emerged and died due to the inhibitory effect of floating naphthalene in water.

In this study, it appeared that all parameters except salt supported the mortality of *Culex* mosquito larvae more than in *Anopheles* mosquito larvae when exposed to powder of *Psoralea corylitolia*. Similarly, all parameters supported the mortality of both larvae exposed to *Sesamum radiatum* from 30 minutes to the end of the post experimental exposure. Physicochemical parameters of mosquito breeding sites and termitarium show that it affects the abundance and distribution of species as reported in the study of Ojjanwuna et al. (2021) and Akpan et al. (2020). Similar observation was reported in the study of Kipyab et al. (2015) as physical and chemical variables of the environment influenced the distribution of malaria vector in the field. Moreover, Emidi et al. (2017) revealed that high conductivity and saltiness of water encourage the abundance of *Anopheles* larvae. The plant powders may have influenced the salt and conductivity, thus causing variable mortalities. Lethal time for 50% of mosquitoes ranged from 31 to 5858min whereas LT<sub>95</sub> ranged from 1435 to 10218min for the two insecticidal test plants respectively. *Culex* mosquitoes exposed to 1.5g of *Psoralea corylitolia* recorded the lowest lethal time and this followed by *Anopheles* mosquitoes exposed to same concentration of *Sesamum radiatum*. The lethal time recorded

in this study is higher than those reported in other studies where plant materials were tried in other insects (Ojjanwuna et al. 2018; 2021).

## 5. Conclusion

This study forms the baseline for the trial of *Sesamum radiatum* and dust of *Psoralea corylitolia* on the mortality of these mosquitoes. It has demonstrated that these two plants did not cause complete mortality on the mosquito larvae. Proper care is required in the interpretation of the result of this study as oil extraction of these plants could cause complete mortality in the mosquitoes. Optimum concentration of *Sesamum radiatum* on both mosquitoes and *Psoralea corylitolia* on *Culex* mosquitoes suppressed the emergence of larvae. More studies on the gas chromatogram of these plants are required to check for the active components.

## Authors' Contributions

CC and VN designed the study, VN, MG and CC carried out the experiment, collected, analyzed field data, and interpreted the data. CC, MG, ET and VN wrote and reviewed the manuscript. All authors read and approved the final manuscript.

## ORCID

Ojjanwuna Chioma C <http://orcid.org/0000-0003-0702-8382>

Enwemiwe Victor N <http://orcid.org/0000-0002-2116-4448>

Ojo Mercy G <http://orcid.org/0000-0001-6238-9312>

Oborayiruvbe Treasure E <http://orcid.org/0000-0001-7285-0723>

## REFERENCES

- Akpan AU, Ojjanwuna CC, Ubulom PME, Clement AY and Oboho DE, 2020. Effect of physico-chemical parameters on the abundance and diversity of termites and other arthropods in termite mounds in Uyo, Akwa Ibom State, Nigeria. *FUDMA Journal of Sciences* 4(2): 92-100. <https://doi.org/10.33003/fjs-2020-0402-206>
- Alam F, Khan GN, Asad and MHHB, 2018. *Psoralea corylifolia* L: Ethnobotanical, biological, and chemical aspects: A review. *Phytotherapy Research* 32(4): 597-615. <https://doi.org/10.1002/ptr.6006>
- Anonymous, 2014. The Federal Republic of Nigeria. National Malaria Strategic Plan 2014-2020. Roll Back Malaria 2014;1-110. Available at: [www.nmcp.gov.ng](http://www.nmcp.gov.ng) (accessed Feb 2, 2017).
- Ayinde AA, Morakinyo OM and Sridhar MKC, 2020. Repellency and larvicidal activities of *Azadirachta indica* seed oil on *Anopheles gambiae* in Nigeria. *Heliyon* 6(5): e03920. <https://doi.org/10.1016/j.heliyon.2020.e03920>
- Becker N, Dušan P, Marija Z, Clive B, Christine D, Minoo M and Achim K, 2020. Mosquitoes Identification, Ecology and Control, 3<sup>rd</sup> Ed. Springer Nature Switzerland AG. <https://doi.org/10.1007/978-3-030-11623-1>
- CDC, 2020. What is a Mosquito? <https://www.cdc.gov/mosquitoes/about/what-is-a-mosquito.html> Accessed 20 Aug 2022.
- Chancey C, Grinev A, Volkova E and Rios M, 2015. The global Ecology and epidemiology of West Nile Virus. *BioMed Research International* 2015: Article # 376230. <https://doi.org/10.1155/2015/376230>
- Dad O, Attaullah M, Ullah H, Ullah R, Ilahi I, Ahmad S, Ahmad B, Ali L and Zeb S, 2019. Prevalence of malaria and status of *Plasmodium* sp. in Dir Lower, Pakistan. *Journal of Biodiversity and Environmental Sciences* 15(2): 83-87.
- Ekedo CM, Okore OO, Obeagu IA and Okafor CC, 2019. Effect of some petroleum products on *Anopheles gambiae* S.L. larvae in Umudike, Ikwuano LGA Abia state, Nigeria. *International Journal of Mosquito Research* 6 (3): 32–36.
- Emidi B, Kisinza WN, Mmbando BP, Malima R and Mosha FW, 2017. Effect of physicochemical parameters on *Anopheles* and *Culex* mosquito larvae abundance in different breeding sites in a rural setting of Muheza, Tanzania. *Parasites Vectors* 10: 304. <https://doi.org/10.1186/s13071-017-2238-x>
- Enwemiwe VN, Anyaele OO and Ojjanwuna CC, 2021. Evaluating the potentials of randomised integrated control trial on tungiasis in a South-western Nigerian community. *Acta Tropica* 223: 106076. <https://doi.org/10.1016/j.actatropica.2021.106076>
- Enwemiwe VN, Ojjanwuna CC and Anyaele OO, 2020. Assessing the potentials of two local topical ointments as affordable treatment against tungiasis infestation: a self-experimentation in Igbokoda, Nigeria. *Parasite Epidemiology and Control* 11: e00168. <https://doi.org/10.1016/j.parepi.2020.e00168>
- Farag SM, Kamel OM, El-Hassan GMA and Zyaan OH, 2021. Larvicidal and repellent potential of *Sesamum indicum* hull peels extracts against *Culex pipiens* L. (Diptera: Culicidae). *Egyptian Journal of Aquatic Biology & Fisheries* 25(2): 995-1011.
- Hamlet A, Dengela D, Tongren JE, Tadesse FG, Bousema T, Sinka M, Seyoum A, Irish SR, Armistead JS and Churcher T, 2022. The potential impact of *Anopheles stephensi* establishment on the transmission of *Plasmodium falciparum* in Ethiopia and prospective control measures. *BMC Medicine* 20(1): 135. <https://doi.org/10.1186/s12916-022-02324-1>
- Kampen H and Walther D, 2018. Vector Potential of Mosquito Species (Diptera: Culicidae) Occurring in Central Europe. In: Benelli G, Mehlhorn H (eds) *Mosquito-borne Diseases*. Parasitology Research Monographs, vol 10. Springer, [https://doi.org/10.1007/978-3-319-94075-5\\_5](https://doi.org/10.1007/978-3-319-94075-5_5)
- Kimani EW, Vulule JM, Kuria IW and Mugisha F, 2006. Use of insecticide-treated clothes for personal protection against malaria: a community trial. *Malaria Journal* 5: 63. <https://doi.org/10.1186/1475-2875-5-63>.

- Kipyab PC, Khaemba BM, Mwangangi JM and Mbogo CM, 2015. The physicochemical and environmental factor affecting the distribution of *Anopheles merus* along the Kenyan Coast. *Parasite Vectors* 8: 221. <https://doi.org/10.1186/s13071-015-0819-0>
- Lai S, Johansson MA, Yin W, Wardrop NA, van Panhuis WG, Wesolowski A, Kraemer MUG, Bogoch II, Kain D, Findlater A, Choisy M, Huang Z, Mu D, Li Y, He Y, Chen Q, Yang J, Khan K, Tatem AJ and Yu H, 2018. Seasonal and interannual risks of dengue introduction from South-East Asia into China, 2005-2015. *PLOS Neglected Tropical Diseases* 12(11): e0006743. <https://doi.org/10.1371/journal.pntd.0006743>
- Lopez I, 2022. What is Skeeter Syndrome? <https://www.webmd.com/allergies/what-is-skeeter-syndrome>. Accessed 20 Aug 2022.
- Mahrn HA, 2022. Using nanoemulsions of the essential oils of a selection of medicinal plants from Jazan, Saudi Arabia, as a green larvicidal against *Culex pipiens*. *PLoS ONE* 17(5): e0267150. <https://doi.org/10.1371/journal.pone.0267150>
- Nwaneli El, Eguonu I, Ebenebe JC, Osuorah CDI, Ofiaeli OC and Nri-Ezedi CA, 2020. Malaria prevalence and its sociodemographic determinants in febrile children - a hospital-based study in a developing community in South-East Nigeria. *Journal of Preventive Medicine and Hygiene* 61: E173-E180. <https://doi.org/10.15167/2421-4248/jpmh2020.61.2.1350>
- Ojianwuna CC and Enwemiwe VN, 2020a. Efficiency of ginger, pepper fruit and alligator pepper in the control of cowpea bruchid (*Callosobruchus maculatus*) (Fabricius, 1775). *FUVW Trends in Science and Technology Journal* 5(2): 445-450.
- Ojianwuna CC and Enwemiwe VN, 2020b. Monitoring the toxicity of dry wood termites using three plant powders in laboratory culture. *Journal of Biopesticides* 13(2): 150-158.
- Ojianwuna CC and Enwemiwe VN, 2021. Toxicity and repellency of scent leave (*Ocimum gratissimum*), kerosene and naphthalene in single form and in mixed form in termites (*Macrotermes bellicosus*). *Journal of Biopesticides* 14 (1): 41-49.
- Ojianwuna CC and Enwemiwe VN, 2022. Insecticidal effectiveness of naphthalene and its combination with kerosene against the emergence of *Aedes aegypti* in Ika North East, LGA, Delta State, Nigeria. *Parasite Epidemiology and Control* 18: e00259. <https://doi.org/10.1016/j.parepi.2022.e00259>
- Ojianwuna CC and Umoru PA, 2010. Effects of *Cymbopogon citratus* (lemon grass) and *Ocimum suave* (wild basil) applied as mixed and individual powders on the egg laid and emergence of adult *Callosobruchus maculatus* (cowpea bruchids). *African Journal of Agricultural Research* 5(20): 2837-2840.
- Ojianwuna CC and Umoru PA, 2011. Acute toxicity of Lemon grass (*Cymbopogon citratus*) and wild basil (*Ocimum suave*) applied as mixed and individual powders against the cowpea bruchids, *Callosobruchus maculatus*, in cowpea. *African Journal of Agricultural Research* 6(29): 6311-6319. <https://doi.org/10.5897/AJAR10.323>
- Ojianwuna CC, Edafemakor AG and Iloh AC, 2011. Toxicity of *Ocimum suave* (wild basil) leaf oil on adult housefly (*Musca domestica*). *International Research Journal of Agricultural Science and Soil Science* 1(10): 417-420.
- Ojianwuna CC, Enwemiwe VN and Erhunmwun S, 2021b. Effect of petroleum products on larvicidal activity of *Aedes* mosquitoes in Ika north-east LGA, Delta state, Nigeria. *Open Ecology Journal* 14: 24-30. <https://doi.org/10.2174/2590277602114010024>
- Ojianwuna CC, Enwemiwe VN and Ossai PI, 2021a. The use of deltamethrin, permethrin, their combination and synergist on female *Anopheles* mosquito in Isoko south, Delta state. *Nigerian Journal of Science and Environment* 19(1): 39-50.
- Ojianwuna CC, Enwemiwe VN, and Ekeazu CN, 2021. Abundance and distribution of *Anopheles* mosquito in relation to physicochemical properties in Delta State, Nigeria. *FUDMA Journal of Sciences* 5(3): 274-280. <https://doi.org/10.33003/fjs-2021-0503-752>
- Ojianwuna CC, Ilondu EM and Akpan AU, 2018. Insecticidal potency OF *Ocimum suave* (wild basil) leave oil extract against maize weevils (*Sitophilus zeamais* Motschulsky) and test of maize seeds' viability. *World Journal of Applied Science and Technology* 10(2): 1-6.
- Ojianwuna CC, Ilondu EM and Enwemiwe VN, 2021. Larvicidal efficacy of leaf extracts from three asteraceous plant against mosquito (*Culex quinquefasciatus*). *FUDMA Journal of Sciences* 5(2): 100-108. <https://doi.org/10.33003/fjs-2021-0502-528>
- Steinbrink A, Brugger K, Margos G, Kraiczy P and Klimpel S, 2022. The evolving story of *Borrelia burgdorferi* sensu lato transmission in Europe. *Parasitology Research* 121(3): 781-803. <https://doi.org/10.1007/s00436-022-07445-3>
- Trájer AJ and Padisák J, 2019. Exploration of the main types of biome-scale culicid entomofauna (Diptera: Culicidae) in Europe and its relationship to the occurrence of mosquito-borne arboviruses. *Acta Zoologica Academiae Scientiarum Hungaricae* 65(3): 299-322. <https://doi.org/10.17109/AZH.65.3.299.2019>
- WHO, 2017. Vector-borne diseases. World Health Organization, Geneva. <https://www.who.int/en/news-room/factsheets/detail/vector-borne-diseases> Assessed August 9, 2022.
- WHO, 2021. World malaria report 2021. World Health Organization, Geneva.